

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
5 February 2004 (05.02.2004)

PCT

(10) International Publication Number  
**WO 2004/011457 A1**

(51) International Patent Classification<sup>7</sup>: C07D 405/12

(21) International Application Number:  
PCT/KR2003/001502

(22) International Filing Date: 26 July 2003 (26.07.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
10-2002-0044164 26 July 2002 (26.07.2002) KR  
10-2003-0013889 6 March 2003 (06.03.2003) KR

(71) Applicants (*for all designated States except US*): YUHAN CORPORATION [KR/KR]; 49-6, Taebang-dong, Tongjak-gu, Seoul 156-020 (KR). DONG-A PHARMACEUTICAL CO., LTD. [KR/KR]; 252, Yongdoo2-dong, Dongdaemoon-gu, Seoul 130-070 (KR).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): LEE, Jong-Wook [KR/KR]; 50-5, Byulyang-dong, Kwacheon-city, Kyunggi-do 427-040 (KR). LEE, Bong-Yong [KR/KR]; 116-2105 Haniltown, Jowon-dong, Jangan-gu, Suwon-city, Kyunggi-do 440-709 (KR). LEE, Chun-Ho [KR/KR]; 101-707 Samho Apt. 471, Gaebong-dong, Guro-gu, Seoul 152-090 (KR). HUR, Yun [KR/KR]; 101-106 Dae-woo Apt., Pyeongchon-dong, Dongan-gu, Anyang-city, Kyunggi-do 431-817 (KR). HAN, Tae-Dong [KR/KR]; B1, 32-28, Ojeon-dong, Uiwang-city, Kyunggi-do 437-817 (KR). KO, Hyun-Kyoung [KR/KR]; Ga-513 Jeil Apt., Mohyeon-dong, Iksan-city, Jeollabuk-do 570-090 (KR). YUN, Suk-Won [KR/KR]; 207-204 Hyosung Apt. 291, Geoyeo-dong, Songpa-gu, Seoul 138-110 (KR). SHIM, Jae-Young [KR/KR]; 112-207 Woosung 6cha

Apt., Sam-dong, Uiwang-city, Kyunggi-do 437-040 (KR). LIM, Joong-In [KR/KR]; 105-1006 LG Village Apt., Mangpo-dong, Paldal-gu, Suwon-city, Kyunggi-do 442-706 (KR). SON, Moon-Ho [KR/KR]; 101-1006 Shinmijoo Apt., Inkye-dong, Paldal-gu, Suwon-city, Kyunggi-do 442-828 (KR). YANG, Jae-Sung [KR/KR]; B-309 Kyungbuk Apt., Nonhyun-dong, Gangnam-gu, Seoul 135-833 (KR). KIM, Mi-Kyung [KR/KR]; 223-1002 Hwanggol Byucksan Apt., Youngtong-dong, Paldal-gu, Suwon-city, Kyunggi-do 442-740 (KR).

(74) Agent: LEE, Young-Pil; The Cheonghwa Building, 1571-18, Seocho-dong, Seocho-gu, Seoul 137-874 (KR).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: 1-PHENYLPYPERIDIN-3-ONE DERIVATIVES AND PROCESSES FOR THE PREPARATION THEREOF

(57) Abstract: The present invention provides 1-phenylpiperidin-3-one derivatives and pharmaceutically acceptable saltsthereof, having cysteine protease inhibitory activity, pharmaceutical compositions containing the same as an active ingredient, and processes for the preparation thereof.

WO 2004/011457 A1

## 1-PHENYLPYPERIDIN-3-ONE DERIVATIVES AND PROCESSES FOR THE PREPARATION THEREOF

### Technical Field

5

The present invention relates to novel 1-phenylpyperidin-3-one derivatives and pharmaceutically acceptable salts thereof having cysteine protease inhibitory activity, pharmaceutical compositions containing the same as an active ingredient, and processes for the preparation thereof.

10

### Background Art

Cysteine proteases, in particular cathepsins of the papain family, participate in the normal physiological protein degradation, such as connective  
15 tissue degradation in animals, including humans. However, elevated levels of these enzymes in the body may cause various diseases. For example, it is reported that calpain protease, one of cysteine proteases, is involved in stroke and neurodegenerative diseases, such as Alzheimer's disease [G. J. Wells, et. al., *Exp. Opin. Ther. Patents*, 8(12), 1707 (1998)]; cathepsin B is involved in  
20 cancer metastasis [S. Michaud, et. al., *Exp. Opin. Ther. Patents*, 8(6), 645 (1998)]; and cathepsin L is involved in chronic rheumatoid arthritis and osteoarthritis [H-H Otto & T. Schirmeister, *Chem. Rev.*, 97, 133 (1997)]. International Publication Nos. WO 00/49007, WO 00/49008 and 00/48992 disclose that cathepsin S is involved in chronic obstructive pulmonary disease  
25 (COPD).

As cysteine proteases other than the papain family, caspase proteases are also known to be involved in inflammatory diseases such as osteoarthritis [D. D. Fairlie et al, *J. Med. Chem.* 43(3), 305 (2000)] and Rhinovirus  
30 3C protease is involved in influenza [Q. M. Wang, *Exp. Opin. Ther. Patents*, 8(9), 151 (1998)].

Moreover, it is known that cathepsin K is distributed selectively in osteoblasts associated with bone resorption in a bone remodeling process and

plays an important role in degradation of organic matter in bones. Accordingly, various attempts have been made to develop a novel agent for treating osteoporosis by inhibiting such proteases [W. W. Smith, et. al., *Exp. Opin. Ther. Patents*, 9(6), 683 (1999); W. Kim, et. al., *Exp. Opin. Ther. Patent*, 12(3), 419 (2002)].

For example, it has been reported that vinylsulfone group-containing peptidomimetic compounds irreversibly inhibit cysteine proteases including cathepsin K [J. T. Palmer, et. al., *J. Med. Chem.*, 38, 3139 (1995); W. W. Roush, et. al., *J. Am. Chem. Soc.*, 120, 10994 (1998)]. Various peptidyl or peptidomimetic compounds having 1,3-diamino-1,3-propan-2-one as a main chain have been also reported to inhibit cysteine proteases including cathepsin K [D. S. Yamashita, et. al., *J. Am. Chem. Soc.*, 119, 11351 (1997); S. K. Thompson, et. al., *Proc. Nacl. Acad. Sci.*, 94, 14249(1997); WO 98/08802; WO 98/48799; WO 98/49152; WO 98/50342; WO 98/50534; and WO 99/11637]. Peptidyl aldehyde (Japanese Patent Publication Nos. 8/092193, 8/151394 and 10/147564, and WO 98/25899) and peptidyl epoxysuccinamides (WO 98/47887) are proposed as irreversible cysteine protease inhibitors. WO 98/50533 proposes peptidyl 3-keto-heterocyclic derivatives, as a cathepsin K inhibitor. Further, it has been also reported that peptidyl  $\alpha$ -ketoamides (EP 10085920); peptidyl phenylethylamines (WO 00/48993); peptidyl cycloketones (WO98/05336); and peptidyl azepines (WO 00/38687 and WO 01/34565) inhibit cysteine proteases, including cathepsin K.

However, those peptidyl compounds as known in the art readily subject to attack of proteolytic enzymes in the body, and therefore, it is difficult to develop drugs based on peptidyl compounds for treating diseases such as osteoporosis [Rasnick, D. *Perspect. Drug Discov. Design*, 6, 47(1996); M. Sato, et. al., *J. Med. Chem.*, 42, 3(1999); W. W. Smith, et. al., *Exp. Opin. Ther. Patents*, 9(6), 683 (1999)]. Further, structural similarities in active sites of cathepsins B and L; and cathepsins K and S make it difficult to develop selective cysteine protease inhibitors each having a satisfactory selectivity to respective cathepsins, such as cathepsin B, L, K, and S [W. Kim, et. al., *Exp. Opin. Ther. Patents*, 12(3), 419 (2002)]. Besides, there has been a demand in

the art to develop cysteine protease inhibitors having sufficient bioavailability and physicochemical stability enabling practical pharmacokinetic evaluations in oral administration.

5 Disclosure of the Invention

The present invention provides a novel compound (i.e., 1-phenylpiperidin-3-one derivative) which exhibits an excellent inhibitory activity against cysteine protease, a high selective inhibition activity against cathepsin K,  
10 and a satisfactory bioavailability. Further, the present invention provides a process for preparing the 1-phenylpiperidin-3-one derivative and a pharmaceutical composition comprising the same.

In one aspect of the present invention, there is provided a 1-phenylpiperidin-3-one derivative or a pharmaceutically acceptable salt  
15 thereof.

In another aspect of the present invention, there is provided a process for preparing the 1-phenylpiperidin-3-one derivative or a pharmaceutically acceptable salt thereof, which comprises oxidizing a synthetic intermediate thereof.

20 In still another aspect of the present invention, there is provided an intermediate for the synthesis of the 1-phenylpiperidin-3-one derivative or a pharmaceutically acceptable salt thereof.


In still another aspect of the present invention, there is provided a process for preparing the synthetic intermediate.

25 In still another aspect of the present invention, there is provided a pharmaceutical composition for inhibiting cysteine proteases comprising a therapeutically effective amount of a 1-phenylpiperidin-3-one derivative or a pharmaceutically acceptable salt thereof as an active ingredient and a pharmaceutically acceptable carrier.

30

Best mode for carrying out the Invention

In accordance with one aspect of the present invention, there is provided a compound of formula (I) or a pharmaceutically acceptable salt thereof:




5 wherein:


$R^1$  is a  $C_{1-6}$  alkyl group optionally substituted with phenyl,  $C_{1-6}$  alkoxy, or benzyloxy; a  $C_{2-6}$  alkenyl group optionally substituted with phenyl; a  $C_{3-6}$  cycloalkyl group; a  $C_{1-5}$  alkoxy group; a phenyl group substituted with halogen, phenyl, trifluoromethoxy, oxopyrrolidyl, mono- or di-  $C_{1-4}$  alkylamino or  $R^4$ - $C_{1-4}$  alkoxy (wherein,  $R^4$  is morpholine, pyrrolidine or piperidine); a furanyl group optionally substituted with one or more functional groups selected from the group consisting of  $C_{1-6}$  alkyl, halogen, and oxopyrrolidyl; a benzofuranyl group optionally substituted with  $C_{1-6}$  alkyl or  $R^4$ - $C_{1-4}$  alkoxy (wherein,  $R^4$  is morpholine, pyrrolidine or piperidine); a thiophenyl group substituted with  $C_{1-6}$  alkyl or halogen; a  $C_{1-6}$  alkyl-isoxazolyl group; a pyridyl group optionally substituted with halogen; a morpholinyl group; a benzothiophenyl group; a quinolinyl group; a pyrazinyl group; a benzyloxy group; an oxopyranyl group; a  $C_{1-6}$  alkyl-7H-imidazo[2,1-b]oxazolyl group; a  $C_{1-6}$  alkyl-chromon-2-yl group; or a (N-t-butoxycarbonyl)piperidinyl group, and

15


20  $R^2$  and  $R^3$  are, each independently, hydrogen; hydroxy; nitro; halogen; cyano; a  $C_{1-6}$  alkyl group optionally substituted with one or more halogen atoms;  $C_{1-5}$  alkoxy;  $C_{1-5}$  alkyl-thio; furyl; 1H-tetrazol-5-yl; oxazolyl; or a group selected from the formula consisting of (II), (III) and (IV):



(II)



(III)



(IV)

wherein,

R<sup>5</sup> is hydrogen; hydroxy; C<sub>1-6</sub> alkyl; C<sub>1-5</sub> alkoxy; mono- or di- C<sub>1-6</sub> alkylamino; or C<sub>3-6</sub> cycloalkylamino,

R<sup>6</sup> is C<sub>1-6</sub> alkyl; phenyl optionally substituted with a C<sub>1-4</sub> alkoxy group; benzyl optionally substituted with a C<sub>1-4</sub> alkoxy group,

R<sup>7</sup> and R<sup>8</sup> are, each independently, hydrogen; a C<sub>1-6</sub> alkylcarbonyl group optionally substituted with halogen, C<sub>1-4</sub> alkoxy, or phenyl; C<sub>2-6</sub> alkenylcarbonyl; C<sub>1-4</sub> alkoxycarbonyl; C<sub>3-6</sub> cycloalkylcarbonyl; benzoyl optionally substituted with one or more halogen atoms; mono- or di- C<sub>1-4</sub> alkylcarbonyl; or C<sub>1-4</sub> alkylsulfonyl, or bonded each ether to form a morpholine, azetidin-2-one, 3,3-dimethylazetidin-2-one, pyrrolidin-2-one, pyrrole, 2,5-dihydropyrrole, piperidin-2-one, oxazolidin-2-one, imidazolidin-2-one, imidazolidin-2,5-dione, tetrazole, 1,1-dioxoisothiazolidine, or C<sub>1-6</sub> alkyl-aziridin-2-one ring, and

n is 0, 1, or 2.

Among the compounds of the present invention, preferred are those wherein: R<sup>1</sup> is a furanyl group optionally substituted with C<sub>1-6</sub> alkyl, halogen, or oxopyrrolidyl; or a benzofuranyl group optionally substituted with C<sub>1-6</sub> alkyl or R<sup>4</sup>-C<sub>1-4</sub> alkoxy (wherein, R<sup>4</sup> is morpholine, pyrrolidine or piperidine) and R<sup>2</sup> and R<sup>3</sup> are, each independently, hydrogen; halogen; or a group of formula (III) or (IV).

More preferred compounds of formula (I) are listed below:

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-nitrophenyl)piperidin-3-one;

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-nitrophenyl)piperidin-3-one;

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(5-chloro-2-nitrophenyl)piperidin-3-one;

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-methanesulfonylphenyl)piperidin-3-one;

- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-acetylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(4-cyanophenyl)piperidin-3-one;
- 5 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-cyano-3-fluorophenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-cyano-6-fluorophenyl)piperidin-3-one;
- 10 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-cyano-5-fluorophenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(3-cyano-5-fluorophenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-ethanesulfonylphenyl)piperidin-3-one;
- 15 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(4-acetylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(4-formylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
20 1-(2-formylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(4-nitrophenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(3-methyl-4-nitrophenyl)piperidin-3-one;
- 25 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(3-fluoro-2-formylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(3-chloro-2-formylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
30 1-(2-formyl-4-nitrophenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-formyl-6-fluorophenyl)piperidin-3-one;

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-fluoro-3-formylphenyl)piperidin-3-one;

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-isopropylsulfonylphenyl)piperidin-3-one;

5 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(4-fluoro-2-trifluoromethylphenyl)piperidin-3-one;

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-cyanophenyl)piperidin-3-one;

10 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-(4-morpholino)phenyl]piperidin-3-one;

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-ethoxyphenyl)piperidin-3-one;

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-fluoro-4-trifluoromethylphenyl)piperidin-3-one;

15 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-chloro-4-trifluoromethylphenyl)piperidin-3-one;

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2,4-dichlorophenyl)piperidin-3-one;

20 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(4-chloro-2-fluorophenyl)piperidin-3-one;

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-methoxy-5-cyanophenyl)piperidin-3-one;

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(4-cyano-2-fluorophenyl)piperidin-3-one;

25 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-(2-furyl)phenyl]piperidin-3-one;

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-bromophenyl)piperidin-3-one;

30 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-methylthiophenyl)piperidin-3-one;

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-(tert-butoxycarbonyl)phenyl]piperidin-3-one;



- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-ethoxycarbonylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-methoxyphenyl)piperidin-3-one;
- 5 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(4-fluoro-2-methylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-isopropoxyphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
10 1-(2-ethylthiophenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-(3,3-dimethylazetidin-2-one-1-yl)phenyl]piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-(pyrrolidin-2-one-1-yl)phenyl]piperidin-3-one;
- 15 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-(piperidin-2-one-1-yl)phenyl]piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-(2-oxazolidinon-1-yl)phenyl]piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
20 1-[2-(2-imidazolidinon-1-yl)phenyl]piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-[N-(ethoxycarbonyl)amino]phenyl]piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-[N-(methoxycarbonyl)amino]phenyl]piperidin-3-one;
- 25 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-[N-(4-chlorobutyryl)amino]phenyl]piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-[N-(isobutyryl)amino]phenyl]piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
30 1-[2-[N-(methoxyacetyl)amino]phenyl]piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-[N-(cyclopropanecarbonyl)amino]phenyl]piperidin-3-one;

- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-[N-(3-phenylpropionyl)amino]phenyl]piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-[N-(2-fluorobenzoyl)amino]phenyl]piperidin-3-one;
- 5 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-[N-(acetyl)amino]phenyl]piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-(2,5-dihydropyrrol-1-yl)phenyl]piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
10 1-[2-(tetrazol-1-yl)phenyl]piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-(pyrrol-1-yl)-5-fluorophenyl]piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-(pyrrol-1-yl)phenyl]piperidin-3-one;
- 15 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-(N,N-dimethylcarbamoyl)phenyl]piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-(N-methylcarbamoyl)phenyl]piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
20 1-(2-carboxyphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(4-carboxyphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-(1,3-oxazol-5-yl)phenyl]piperidin-3-one;
- 25 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-hydroxyphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-(1H-tetrazol-5-yl)phenyl]piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
30 1-(2-methylsulfinylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-ethylsulfinylphenyl)piperidin-3-one;

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-isopropylsulfinylphenyl)piperidin-3-one;

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-N-(acryloyl)aminophenyl]piperidin-3-one;

5 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-(3,3-dimethylureido)phenyl]piperidin-3-one;

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-(methoxycarbonyl)phenyl]piperidin-3-one;

10 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-phenylpiperidin-3-one;

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-(1,1-dioxo-isothiazolin-2-yl)phenyl]piperidin-3-one;

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-met  
hanesulfonylphenyl)piperidin-3-one;

15 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(3-fluo  
ro-2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(4-fluo  
ro-2-nitrophenyl)piperidin-3-one;

20 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(4-fluo  
ro-2-formylphenyl)piperidin-3-one;

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-isop  
ropylsulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-p  
henylsulfonylphenyl)piperidin-3-one;

25 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-(4  
-methoxybenzylsulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(4-fluo  
ro-2-methanesulfonylphenyl)piperidin-3-one;

30 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(4-met  
hoxy-2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(4-fluo  
ro-2-thiomethoxyphenyl)piperidin-3-one;

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-cyanophenyl)piperidin-3-one;

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(4-fluoro-2-methoxyacetylaminophenyl)piperidin-3-one;

5 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(4-fluoro-2-methoxycarbonylaminophenyl)piperidin-3-one;

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(4-fluoro-2-cyclopropylcarbonylaminophenyl)piperidin-3-one;

10 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(4-fluoro-2-isobutyrylaminophenyl)piperidin-3-one;

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[4-fluoro-2-(4-chlorobutyryl)aminophenyl]piperidin-3-one;

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[4-fluoro-2-(3,3-dimethylazetidin-2-one-1-yl)phenyl]piperidin-3-one;

15 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(3-methyl-aziridin-2-one-1-yl)phenyl]piperidin-3-one;

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(azetidin-2-one-1-yl)phenyl]piperidin-3-one;

20 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[4-fluoro-2-(pyrrolidin-2-one-1-yl)phenyl]piperidin-3-one;

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(1,3-imidazolidin-2,5-dione-1-yl)phenyl]piperidin-3-one;

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[4-fluoro-2-(1,1-dioxo-isothiazolidin-2-yl)phenyl]piperidin-3-one;

25 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[4-fluoro-2-(piperidin-2-one-1-yl)phenyl]piperidin-3-one;

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[4-fluoro-2-(pyrrol-1-yl)]piperidin-3-one;

30 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[4-fluoro-2-(oxazol-4-yl)phenyl]piperidin-3-one;

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[4-fluoro-2-(N-methylcarbamoyl)phenyl]piperidin-3-one;

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[2-(N,N-dimethylcarbamoyl)phenyl]piperidin-3-one;

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[2-(N-cyclopropylcarbamoyl)phenyl]piperidin-3-one;

5 4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-aminophenyl)piperidin-3-one;

4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-nitrophenyl)piperidin-3-one;

10 4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-ethanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[4-fluoro-2-(3,3-dimethyl-azetidin-2-one-1-yl)phenyl]piperidin-3-one;

15 4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[4-fluoro-2-(pyrrolidin-2-one-1-yl)phenyl]piperidin-3-one;

4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[4-fluoro-2-(oxazolidin-2-one-3-yl)phenyl]piperidin-3-one;

20 4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[4-fluoro-2-(imidazolidin-2-one-1-yl)phenyl]piperidin-3-one;

4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[2-(3,3-dimethyl-azetidin-2-one-1-yl)phenyl]piperidin-3-one;

4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[2-(pyrrolidin-2-one-1-yl)phenyl]piperidin-3-one;

25 4-[N-[1-[N-(5-chloro-thiophen-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(5-chloro-thiophen-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-ethanesulfonylphenyl)piperidin-3-one;

30 4-[N-[1-[N-(5-chloro-thiophen-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-isopropanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(5-chloro-thiophen-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(5-fluoro-2-methanesulfonylphenyl)piperidin-3-one;

- 4-[N-[1-[N-(5-methyl-thiophen-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(5-methyl-thiophen-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-isopropanesulfonylphenyl)piperidin-3-one;
- 5 4-[N-[1-[N-(5-methyl-thiophen-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(5-fluoro-2-methanesulfonylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(3-methyl-thiophen-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(2-methyl-furan-3-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;
- 10 4-[N-[1-[N-(5-methyl-furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(5-bromo-furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonyl-4-methoxyphenyl)piperidin-3-one;
- 15 4-[N-[1-[N-(4,5-dimethyl-furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(4,5-dimethyl-furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[2-(pyrrolidin-2-one-1-yl)phenyl]piperidin-3-one;
- 4-[N-[1-[N-(morpholin-4-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;
- 20 4-[N-[1-[N-(benzothiophen-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzothiophen-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-tert-butoxycarbonylphenyl)piperidin-3-one;
- 25 4-[N-[1-[N-(benzothiophen-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-chloro-5-trifluoromethylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzothiophen-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(4-chloro-2-fluoro-phenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzothiophen-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methylphenyl)piperidin-3-one;
- 30 4-[N-[1-[N-(3-methyl-benzofuran-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

- 4-[N-[1-[N-(3-methyl-benzofuran-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(5-fluoro-2-trifluoromethylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(3-methyl-benzofuran-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(5-cyano-2-methoxyphenyl)piperidin-3-one;
- 5 4-[N-[1-[N-(quinolin-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(quinolin-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[2-(pyrrolidin-2-one-1-yl)phenyl]piperidin-3-one;
- 4-[N-[1-[N-(quinolin-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-nitrophenyl)piperidin-3-one;
- 10 4-[N-[1-[N-(pyrazin-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(pyrazin-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one dihydrochloride;
- 15 4-[N-[1-[N-(isonicotinylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(6-chloronicotinylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(6-chloronicotinylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(4-fluoro-2-nitrophenyl)piperidin-3-one;
- 20 4-[N-[1-[N-(pyridin-2-carbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(pyridin-2-carbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one hydrochloride;
- 25 4-[N-[1-[N-(4-fluorobenzoyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(4-fluorobenzoyl)amino]cyclohexylcarbonyl]amino]-1-(4-fluoro-2-methanesulfonylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(4-fluorobenzoyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonyl-4-methoxyphenyl)piperidin-3-one;
- 30 4-[N-[1-[N-(4-fluorobenzoyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-trifluoromethylphenyl)piperidin-3-one;

4-[N-[1-[N-(4-fluorobenzoyl)amino]cyclohexanecarbonyl]amino]-1-[(2-(3-phenylpropionylamino)phenyl)piperidin-3-one];

4-[N-[1-[N-(4-fluorobenzoyl)amino]cyclohexanecarbonyl]amino]-1-[2-(cyclopropylamino)phenyl]piperidin-3-one;

5 4-[N-[1-[N-(4-fluorobenzoyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(2,3,4-trifluorobenzoylamino)phenyl]piperidin-3-one;

4-[N-[1-[N-(4-biphenylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

10 4-[N-[1-[N-(4-trifluoromethylbenzoyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(4-(2-morpholinethoxy)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(4-(2-pyrrolidinethoxy)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

15 4-[N-[1-[N-(4-(2-piperidinethoxy)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(4-(2-morpholinethoxy)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one hydrochloride;

20 4-[N-[1-[N-(4-(2-pyrrolidinethoxy)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one hydrochloride;

4-[N-[1-[N-(4-(2-piperidinethoxy)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one hydrochloride;

4-[N-[1-[N-(5-(2-morpholinethoxy)benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

25 4-[N-[1-[N-(5-(2-pyrrolidinethoxy)benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(5-(2-piperidinethoxy)benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

30 4-[N-[1-[N-(5-(2-morpholinethoxy)benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one hydrochloride;



4-[N-[1-[N-(5-(2-pyrrolidinethoxy)benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one hydrochloride;

5 4-[N-[1-[N-(5-(2-piperidinethoxy)benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one hydrochloride;

4-[N-[1-[N-(4-(2-oxopyrrolidine)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(4-(2-oxopyrrolidine)furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

10 4-[N-[1-[N-(4-(2-morpholinethoxy)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-methanesulfonylaminophenyl)piperidin-3-one;

4-[N-[1-[N-(4-(2-morpholinethoxy)benzoyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(pyrrolidin-2-one-1-yl)phenyl]piperidin-3-one;

15 4-[N-[1-[N-(4-(dimethylamino)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(4-(diethylamino)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(4-(dimethylamino)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-methanesulfonylphenyl)piperidin-3-one hydrochloride;

20 4-[N-[1-[N-(4-(diethylamino)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-methanesulfonylphenyl)piperidin-3-one hydrochloride;

4-[N-[1-[N-(3-phenylpropionyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

25 4-[N-[1-[N-(2-methylcinnamoyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(benzyloxycarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-ethylamino)piperidin-3-one;

4-[N-[1-[N-(benzyloxycarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-trifluoromethylphenyl)piperidin-3-one;

30 4-[N-[1-[N-(2-pyran-2-one-5-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-((6-methyl)-7H-imidazo[2,1-b]oxazol-5-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-((6-methyl)-chromon-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

5 4-[N-[1-[N-(isobutyryl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(isovaleryl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

10 4-[N-[1-[N-(3,3-dimethylacryloyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(2-methoxyacetyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(cyclopropanecarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

15 4-[N-[1-[N-(cyclobutanecarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(cyclopentanecarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

20 4-[N-[1-[N-(cyclohexanecarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-((3-isobutyloxy)propionyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-((3-benzyloxy)propionyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;


25 4-[N-[1-[N-((N-t-butoxycarbonyl)piperidin-4-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one.

Furthermore, the present invention encompasses within its scope those pharmaceutically acceptable salts of the compounds of formula (I). The pharmaceutically acceptable salts which fall within the scope of the present invention include organic acid salts or inorganic acid salts, such as hydrochloride, hydrobromide, sulfate, sulfarmate, phosphate, nitrate, acetate, propionate,

succinate, glycolate, stearate, malate, hydroxymalate, phenylacetate, glutamate, benzoate, salicylate, sulfanylate, 2-acetoxy-benzoate, fumarate, toluenesulfonate, methanedisulfonate, ethanedisulfonate, oxalate, or trifluoroacetate.

In accordance with another aspect of the present invention, there is provided a process for preparing the compound of formula (I) or its pharmaceutically acceptable salt, which comprises oxidizing a compound of formula (V) with an oxidizing agent, in accordance with the following Reaction Scheme 1 described below:

Reaction Scheme 1




In the above Reaction Scheme 1,  $R^1$ ,  $R^2$ , and  $R^3$  are the same as defined above.

The oxidizing agent includes, but not limited to, a pyridine- $\text{SO}_3$  complex, oxalyl chloride-dimethylsulfoxide (Swern oxidation), and pyridinium chlorochromate (Jones oxidation).

The reaction may be performed in the presence or absence of a base. The base used includes, but not limited to, triethylamine, N,N-diisopropylethylamine and N-methylmorpholine. And, the reaction may be performed preferably in the presence of an organic solvent, including

dichloromethane, tetrahydrofuran, dimethylformamide, or dimethylsulfoxide. Further, the reaction may be carried out at  $-78^{\circ}\text{C} \sim 50^{\circ}\text{C}$ .

In accordance with still another aspect of the present invention, there is provided an intermediate for the synthesis of the compound of formula (I) or a pharmaceutically acceptable salt thereof. That is, there is provided a compound of formula (V) or a pharmaceutically acceptable salt thereof:




(V)

wherein,  $\text{R}^1$ ,  $\text{R}^2$ , and  $\text{R}^3$  are the same as defined above.

In accordance with still another aspect of the present invention, there is provided a process for preparing the synthetic intermediate, i.e., a compound of formula (V) or its pharmaceutically acceptable salt, which comprises reacting a compound of formula (VI) with a compound of formula (VII), in accordance with the following Reaction Scheme 2 described below:

Reaction Scheme 2



In the above Reaction Scheme 2,  $R^1$ ,  $R^2$ , and  $R^3$  are the same as defined above and X is halogen, nitro,  $C_{1-7}$  alkylsulfonyl, or trifluorosulfonate.

In the above Reaction Scheme 2, the compound of formula (VI) may be prepared using a known method [see, e.g., *J. Med. Chem.* 1994, 30, 999-1014].

5        The reaction may be performed preferably in the presence of a base, including organic bases, such as triethylamine, N,N-diisopropylethylamine, and N-methylmorpholine; and inorganic bases, such as sodium bicarbonate, calcium carbonate, and potassium carbonate. And, the reaction may be performed preferably in the presence of an organic solvent, including  
10    tetrahydrofuran, methanol, ethanol, butanol, acetonitrile, dimethylacetamide, dimethylformamide, or dimethylsulfoxide. Further, the reaction may be carried out at  $0^\circ\text{C} \sim 150^\circ\text{C}$ , preferably at  $50^\circ\text{C} \sim 120^\circ\text{C}$ .

Moreover, the reaction according to Reaction Scheme 2 may be performed preferably in the presence of a palladium catalyst, such as palladium  
15    diacetate ( $\text{Pd}(\text{OAc})_2$ ), and tris(dibenzylideneacetone)dipalladium ( $\text{Pd}_2(\text{dba})_2$ ), to facilitate the reaction by reducing reaction time and increase the yield of product. In case of carrying out the reaction in the presence of the palladium catalyst, the process may be performed preferably in the presence of a ligand and/or a base additionally. Example of the ligand includes  
20    (S)-2,2-bis(diphenylphosphino)-1,1'-binaphthyl (BINAP), 1,1'-bis(diphenylphosphino)ferrocene (dppf), and (tri-O-tolyl)phosphine ( $\text{P}(\text{O-Tol})_3$ ) and example of the base includes cesium carbonate ( $\text{CsCO}_3$ ), sodium t-butoxide ( $\text{t-BuONa}$ ), and potassium t-butoxide ( $\text{t-BuOK}$ ). Further, in case of carrying out the reaction in the presence of the palladium catalyst, the  
25    process may be carried out in the presence of a solvent, such as benzene, toluene, dioxane, or tetrahydrofuran and at  $50^\circ\text{C} \sim 150^\circ\text{C}$ , more preferably at  $80^\circ\text{C} \sim 110^\circ\text{C}$ .

In accordance with still another aspect of the present invention, there is provided a pharmaceutical composition for inhibiting cysteine proteases  
30    comprising a therapeutically effective amount of a 1-phenylpiperidin-3-one derivative or a pharmaceutically acceptable salt thereof as an active ingredient and a pharmaceutically acceptable carrier.

The pharmaceutically acceptable carrier may be a conventional liquid or solid carrier and the compositions of the present invention may further include an excipient (e.g. lactose, sucrose, corn starch, etc.), lubricating agent (e.g. magnesium stearate, etc.), emulsifier, suspending agent, buffering agent and isotonic agent, sweetener and/or flavoring agent, if necessary.

The compositions of the present invention may be administered, either orally or parenterally. The dosage form for the pharmaceutical composition includes oral dosage forms such as tablets, capsules, aqueous solvents, emulsions, suspensions, etc., and paenteral dosage forms such as injections (e.g. subcutaneous, intravenous, intramuscular and intraperitoneal injections) and external application forms (e.g. transdermal preparation, ointments, etc.).

The pharmaceutical compositions of the present invention can be administered to mammals, including human, to treat osteoporosis, osteoarthritis, hypercalcemia disease, Pagets disease, rheumatoid arthritis or bone related disease by inhibiting the activity of cysteine protease superfamily. Further, the compositions may be administered to patients suffered from stroke, Alzheimers disease, chronic obstructive pulmonary disease (COPD), influenza, hypertenstion, tumor or cancer metastasis.

The pharmaceutical compositions of the present invention may be administered in a daily dose of about  $0.1\mu\text{g}$  to  $1000\text{ mg/kg}$  body weight, in a single dose or in divided doses. However, it should be understood that the amount of the active ingredient actually administered ought to be determined considering various relevant factors including the condition to be treated, the chosen route of administration, the age and weight of the individual patient, and the severity of the patient's symptom; and, therefore, the above dose should not be intended to limit the scope of the invention in any way.

The following Examples are given for the purpose of illustration only, and are not intended to limit the scope of the invention.

#### Preparation Example 1:

Preparation of tert-butyl 4-amino-hydroxypiperidine-1-carboxylate

Step 1. N-tert-butoxycarbonyl-1,2,3,6-tetrahydropyridine

The solution of 1,2,3,6-tetrahydropyridine (50 g, 0.60 mol) in 500 ml of dichloromethane was cooled to 0 °C and was added di-tert-butoxydicarbonate  
5 (144g, 0.66 mol) thereto. The reaction mixture was stirred for about 2 hours at room temperature and concentrated to give the titled compound as an oily form (110 g).

$$R_f = 0.71 \text{ [n-Hexane/EtOAc (v/v) = 3/1]}$$

10 Step 2: tert-butyl 7-oxo-3-azabicyclo[4.1.0]heptane-3-carboxylate

To the solution of N-tert-butoxycarbonyl-1,2,3,6-tetrahydropyridine (110 g, 0.60 mol) obtained in Step 1 in 300 ml of dichloromethane, was added 3-chloroperbenzoic acid (m-CPBA; 311g, 1.80 mol). The reaction mixture was  
15 stirred overnight at room temperature and filtered to remove benzoic acid produced. The resulting filtrate was washed with a saturated potassium carbonate solution, brine and distilled water. The resulting organic layer was dried and concentrated on magnesium sulfate to give the titled compound (180.6 g).

20  $R_f = 0.62 \text{ (n-Hexane/ EtOAc (v/v) = 3/1)}$

$^1\text{H NMR}$  (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  8.15-7.93 (m, 1H), 7.77-7.51 (d, 1H), 7.49-7.36 (q, 1H), 3.87-3.63 (m, 2H), 3.39-3.13 (m, 2H), 2.12-1.88 (m, 2H), 1.50 (s, 9H)

25 Step 3. tert-Butyl 4-azido-3-hydroxypiperidine-1-carboxylate

To the solution of tert-butyl 7-oxo-3-azabicyclo[4.1.0]heptane-3-carboxylate (180.6 g, 0.60 mol) obtained in Step 2 in a mixed solvent of 700 ml of methanol and 100 ml of water, were  
30 added ammonium chloride ( $\text{NH}_4\text{Cl}$ ; 64.0g, 1.2 mol) and sodium azide ( $\text{NaN}_3$ ; 98.0g, 1.5 mol). The reaction mixture was refluxed for 8 hours and evaporated under reduced pressure to remove methanol. The resulting residue was

extracted with ethyl acetate and then dried on magnesium sulfate. The resulting residue was purified with column chromatography to give the titled compound (62.0 g).

R<sub>f</sub> = 0.43 (n-Hexane/ EtOAc (v/v) = 3/1)

5

#### Step 4. tert-Butyl 4-amino-3-hydroxypiperidine-1-carboxylate

tert-Butyl 4-azido-3-hydroxypiperidine-1-carboxylate (62 g) obtained in Step 3 was dissolved in 300 ml of methanol. The mixture was hydrogenated  
10 with 10%-palladium at room temperature under 35 psi for about 48 hours and filtered. The resulting filtrate was concentrated to give the titled compound (55 g) as a colorless and oily form.

#### Preparation Example 2:

#### 15 Preparation of 1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexane carboxylic acid

##### Step 1. Methyl 1-[N-(benzofuran-2-ylcarbonyl)amino]-1-cyclohexanecarboxylate

The solution of methyl 1-amino-1-cyclohexanecarboxylate (66.1 g, 340  
20 mmol) and benzofuran-2-carboxyl acid (60.0 g) in 500 ml of dichloromethane was cooled to 0 °C and were added dimethylaminopyridine (DMAP, 50.0 g) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC-HCl; 85.0 g) thereto. The reaction mixture was stirred for about 24 hours at room temperature and then ice water (50 g) was added thereto to stop the reaction.  
25 The organic layer was separated and concentrated to obtain a residue, which was then dissolved with 400 ml of ethyl acetate. The resulting solution was washed with 10% citric acid solution and saturated sodium bicarbonate solution, and then dried and concentrated on sodium sulfate to give the titled compound (105.0 g). The titled compound was used in the next-step without further  
30 purifications.

##### Step 2. 1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarboxylic acid



To the solution of methyl  
1-[N-(benzofuran-2-ylcarbonyl)amino]-1-cyclohexanecarboxylate (105.0 g) in  
5 450 ml of methanol, was added 2N-sodium hydroxide solution (170 ml, 0.34  
mol). The reaction mixture was refluxed for 23 hours and concentrated to  
obtain a residue, which was diluted with 400 ml of water and washed with 300  
ml of diethyl ether. 6N-hydrochloride was added dropwise to the resulting  
aqueous layer to adjust the pH of the solution to pH 3. The resulting white  
10 solid was filtered and dried to give the titled compound (49.5 g).

Example 1:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-m  
15 ethylsulfonylphenyl)piperidin-3-one

Step 1.

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-tert-b  
utoxycarbonyl-3-piperidinol

20

To the solution of tert-butyl 4-amino-3-hydroxypiperidine-1-carboxylate  
(20 g, 92 mmol) obtained in step 4 of Preparation Example 1 and  
1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarboxylic acid (26 g, 92  
mmol) obtained in step 2 of Preparation Example 2 in 150 ml of  
25 dimethylformamide, were added hydroxybenzotriazole(HOBt; 24g, 180 mmol)  
and diisopropylethylamine (DIEA; 23 g, 180 mmol). The reaction mixture was  
cooled to 0°C and was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide  
hydrochloride (EDC-HCl; 35 g, 180 mmol) thereto. The reaction mixture was  
stirred for 4 hours at room temperature, diluted with 300 ml of ethyl acetate,  
30 and washed with 10% citric acid solution, a saturated sodium bicarbonate  
solution and brine. The resulting organic layer was dried and concentrated on  
sodium sulfate. The resulting residue was purified with column

chromatography to give 30 g of the titled compound.

Step 2.

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-3-piper  
5 idinol hydrochloride

The solution of  
4-1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonylamino-1-tert-butox  
ycarbonyl-3-piperidinol (30g) obtained in Step 1 in 100ml of  
10 3N-hydrochloride/ethyl acetate solution was stirred for about 2 hours at room  
temperature and was added 100ml of diethyl ether thereto. The resulting  
white solid was filtered and dried to give 25g of the titled compound.

Step 3.

15 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-m  
ethylsulfonylphenyl)piperidin-3-ol

To the solution of 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]  
cyclohexanecarbonyl]amino]-3-piperidinol hydrochloride (10 g, 23.7 mmol)  
20 obtained in Step 2 and 1-fluorophenyl methyl sulfone (4.9 g, 28.4 mmol) in 50  
ml of dimethylformamide, was added potassium carbonate (8.2 g, 59.3 mmol).  
The reaction mixture was stirred overnight at about 100°C, cooled to room  
temperature, diluted with 200ml of ethyl acetate, and washed with brine. The  
resulting organic layer was dried and concentrated on sodium sulfate. The  
25 resulting residue was purified with column chromatography to give 4.9 g of the  
titled compound as white solid.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.03 (m, 1H), 7.69 (d, 1H), 7.62(t, 1H),  
7.51 (m, 2H), 7.42 (t, 1H), 7.31 (d, 1H), 7.25 (m, 2H), 7.05 (bs, 1H), 6.81 (s, 1H),  
30 4.24 (bs, 1H), 3.90 (m, 1H), 3.70 (m, 1H), 3.36 (m, 2H), 3.30 (m, 1H), 3.24 (s,  
3H), 2.90 (m, 1H), 2.67 (t, 1H), 2.27 (t, 2H), 2.05 (m, 3H), 1.75 (m, 3H), 1.52 (m,  
3H), 1.41 (m, 1H)

## Step 4.

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methylsulonylphenyl)piperidin-3-one

5

To the solution of 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methylsulfonylphenyl)piperidin-3-ol (0.49g, 0.91 mmol) obtained in Step 3 in 3 ml of dimethyl sulfoxide (DMSO), was added dropwise 0.5 ml of triethylamine. The reaction mixture was cooled to 0°C.

10 Pyridine-SO<sub>3</sub> complex (0.5 g, 0.31 mmol) was added to the reaction mixture, which was then stirred for 2 hours, diluted with 20 ml of ethyl acetate, and washed with brine. The resulting organic layer was dried and concentrated on sodium sulfate. The resulting residue was purified with column chromatography to give 210 mg of the titled compound as white solid.

15 mp = 219-220 °C

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.07 (t, 1H), 7.68 (m, 2H), 7.55-7.26 (m, 7H), 6.73 (s, 1H), 4.72 (t, 1H), 3.77 (d, 1H), 3.65 (d, 1H), 3.61 (d, 1H), 3.17 (s, 3H), 2.68 (m, 1H), 2.33 (t, 2H), 2.05 (m, 2H), 1.78 (m, 4H), 1.42-1.22 (m, 2H)

Mass (M+H) = 538

20

Example 2:

## Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-nitrophenyl)piperidin-3-one

25

In accordance with the same procedure as in Example 1, except that 2-fluoronitrobenzene was used instead of 1-fluorophenyl methyl sulfone in Step 3 thereof, 170 mg of the titled compound was prepared.

30 <sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.69 (d, 1H), 7.57 (m, 2H), 7.49 (m, 2H), 7.31 (m, 1H), 7.06 (t, 1H), 6.97 (d, 1H), 6.73 (s, 1H), 4.71 (m, 1H), 3.86 (q, 3H), 3.41 (t, 1H), 2.78 (m, 1H), 2.31 (broad, 2H), 2.00 (m, 3H), 1.75 (m, 2H), 1.52 (m,

2H), 1.38 (m, 1H)

Example 3:

Preparation of

5 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-nitrophenyl)piperidin-3-one

10 In accordance with the same procedure as in Example 1, except that 1,4-difluoro-2-nitrobenzene was used instead of 1-fluorophenyl methyl sulfone in Step 3 thereof, 250 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.68 (d, 1H), 7.66 (d, 1H), 7.50 (m, 4H), 7.45 (m, 3H), 6.72 (s, 1H), 4.67 (m, 1H), 3.68 (s, 2H), 3.33 (m, 2H), 2.72 (m, 1H), 2.32 (m, 2H), 2.04 (m, 3H), 1.74 (m, 4H), 1.27 (m, 2H), 1.21 (m, 1H)

15 Example 4:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(5-chloro-2-nitrophenyl)piperidin-3-one

20 In accordance with the same procedure as in Example 1, except that 2,4-dichloronitrobenzene was used instead of 1-fluorophenyl methyl sulfone in Step 3 thereof, 210 mg of the titled compound was prepared.

25 <sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.75 (d, 1H), 7.72 (d, 1H), 7.67 (m, 2H), 7.07 (m, 1H), 6.96 (s, 1H), 6.93 (d, 2H), 6.87 (s, 1H), 4.72 (m, 1H), 3.72 (s, 2H), 3.46 (m, 2H), 2.53 (m, 1H), 2.25 (m, 2H), 2.02 (m, 3H), 1.76 (m, 4H), 1.43 (m, 2H), 1.13 (m, 1H)

Example 5:

Preparation of

30 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-methanesulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 1, except that 4-fluorophenyl methyl sulfone was used instead of 1-fluorophenyl methyl sulfone in Step 3 thereof, 190 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.67 (m, 3H), 7.52 (m, 3H), 7.44 (m, 1H), 6.93 (m, 2H), 6.72 (s, 1H), 4.75 (m, 1H), 3.69 (d, 1H), 3.52 (d, 2H), 3.04 (m, 2H), 2.84 (s, 3H), 2.70 (m, 1H), 2.25 (m, 2H), 2.04 (m, 2H), 1.76 (m, 4H), 1.33 (m, 3H), 1.21 (m, 1H)

#### Example 6:

10 Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-acetylphenyl)piperidin-3-one

15 In accordance with the same procedure as in Example 1, except that 2-fluoroacetophenone was used instead of 1-fluorophenyl methyl sulfone in Step 3 thereof, 110 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.68 (d, 1H), 7.66 (m, 2H), 7.53 (m, 3H), 7.08 (m, 2H), 6.72 (s, 1H), 4.83 (m, 1H), 3.69 (q, 2H), 3.31 (m, 2H), 2.68 (m, 1H), 2.53 (s, 3H), 2.29 (m, 2H), 2.04 (m, 2H), 1.71 (m, 4H), 1.27 (m, 3H), 1.25 (m, 1H)

#### Example 7:

Preparation of

25 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-cyanophenyl)piperidin-3-one

In accordance with the same procedure as in Example 1, except that 4-fluorobenzonitrile was used instead of 1-fluorophenyl methyl sulfone in Step 3 thereof, 120 mg of the titled compound was prepared.

30 <sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.69 (d, 1H), 7.53 (m, 6H), 7.52 (m, 1H), 6.82 (s, 1H), 4.82 (m, 1H), 4.17 (d, 1H), 3.93 (d, 1H), 3.55 (m, 1H), 3.51 (m, 1H), 2.14 (m, 3H), 1.98 (m, 4H), 1.75 (m, 5H), 1.63 (m, 2H), 1.24 (m, 1H)

Example 8:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-cy  
5 ano-3-fluorophenyl)piperidin-3-one

In accordance with the same procedure as in Example 1, except that 2,6-difluorobenzonitrile was used instead of 1-fluorophenyl methyl sulfone in Step 3 thereof, 170 mg of the titled compound was prepared.

10  $^1\text{H-NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  7.69 (d, 1H), 7.53 (m, 6H), 6.82 (s, 1H), 4.69 (m, 1H), 4.13 (d, 1H), 3.90 (d, 1H), 3.81 (m, 1H), 3.57 (m, 1H), 2.83 (m, 1H), 2.02 (m, 2H) 1.98 (m, 4H), 1.77 (m, 4H), 1.68 (m, 2H), 1.26 (m, 1H)

Example 9:

15 Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-cy  
ano-6-fluorophenyl)piperidin-3-one

20 In accordance with the same procedure as in Example 1, except that 2,3-difluorobenzonitrile was used instead of 1-fluorophenyl methyl sulfone in Step 3 thereof, 220 mg of the titled compound was prepared.

$^1\text{H-NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  7.69 (d, 1H), 7.67 (m, 2H), 7.45 (m, 3H), 7.33(m, 1H), 6.74 (s, 1H), 4.87 (m, 1H), 4.03 (d, 1H), 3.82 (d, 1H), 3.73 (m, 1H), 3.52 (m, 1H), 2.87(m, 1H), 2.34 (m, 2H) 1.99 (m, 4H), 1.77 (m, 4H), 1.68 (m,  
25 2H), 1.26 (m, 1H)

Example 10:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-cy  
30 ano-5-fluorophenyl)piperidin-3-one

In accordance with the same procedure as in Example 1, except that

2,4-difluorobenzonitrile was used instead of 1-fluorophenyl methyl sulfone in Step 3 thereof, 210 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.69 (d, 1H), 7.67 (m, 2H), 7.47 (m, 3H), 7.43 (m, 1H), 6.72 (s, 1H), 4.67 (m, 1H), 4.16 (d, 1H), 3.97 (d, 1H), 3.68 (m, 1H), 3.59 (m, 1H), 2.75 (m, 1H), 2.30 (m, 2H) 1.99 (m, 4H), 1.77 (m, 4H), 1.68 (m, 2H), 1.26 (m, 1H)

#### Example 11:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(3-cyano-5-fluorophenyl)piperidin-3-one

In accordance with the same procedure as in Example 1, except that 3,5-difluorobenzonitrile was used instead of 1-fluorophenyl methyl sulfone in Step 3 thereof, 200 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.69 (d, 1H), 7.55 (m, 5H), 7.26 (m, 1H), 6.72 (m, 1H), 4.66 (m, 1H), 4.11 (d, 1H), 3.85 (d, 1H), 3.51 (m, 2H), 2.92 (m, 1H), 2.32 (m, 3H), 1.98 (m, 3H), 1.75 (m, 6H), 1.68 (m, 4H), 1.53(m, 1H)

#### Example 12:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-ethylsulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 1, except that 1-fluorophenyl ethyl sulfone was used instead of 1-fluorophenyl methyl sulfone in Step 3 thereof, 180 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.05 (d, 1H), 7.70 (m, 2H), 7.50 (m, 3H), 7.43 (m, 3H), 6.73 (s, 1H), 4.71 (m, 1H), 3.75 (d, 1H), 3.55 (d, 1H), 3.53 (d, 1H), 3.35 (m, 2H), 3.17 (t, 1H), 2.71 (m, 1H), 2.30 (m, 2H), 2.05 (m, 2H), 1.91 (m, 1H), 1.74 (m, 3H), 1.55 (m, 2H), 1.35 (m, 1H) 1.12 (t, 3H)

Example 13:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-  
ethylphenyl)piperidin-3-one

5

In accordance with the same procedure as in Example 1, except that 4-fluoroacetophenone was used instead of 1-fluorophenyl methyl sulfone in Step 3 thereof, 120 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) : 8.00 (d, 1H), 7.98 (d, 1H), 7.89 (m, 1H), 7.51  
10 (m, 4H), 7.48 (m, 2H), 6.82(s, 1H), 4.87 (m, 1H), 4.21 (d, 1H), 3.95 (d, 1H),  
3.64(t, 1H), 3.16 (m, 1H), 2.46 (s, 3H), 2.15 (m, 3H), 1.98 (m, 5H), 1.76 (m, 6H),  
1.74 (m, 3H)

Example 14:

15 Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-for  
mylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 1, except that  
20 4-fluorobenzadehyde was used instead of 1-fluorophenyl methyl sulfone in Step  
3 thereof, 150 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 9.78 (s, 1H), 7.67(d, 2H), 7.74 (d, 1H),  
7.68 (m, 2H), 7.47 (m, 2H), 7.31 (m, 1H), 6.86 (m, 1H), 6.74 (s, 1H), 4.70 (m,  
1H), 4.22 (d, 1H), 3.99 (d, 1H), 3.95 (m, 1H), 3.60 (m, 2H), 2.87 (m, 1H),  
25 (m, 2H), 1.98 (m, 2H), 1.75 (m, 3H), 1.52 (m, 2H)

Example 15:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-for  
30 mylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 1, except that



2-fluorobenzaldehyde was used instead of 1-fluorophenyl methyl sulfone in Step 3 thereof, 130 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 10.21 (s, 1H), 7.95 (d, 1H), 7.87 (d, 1H), 7.67 (m, 2H), 7.55 (m, 2H), 7.47 (m, 1H), 7.31 (m, 1H), 6.73 (s, 1H), 4.68 (m, 1H), 3.76 (s, 2H), 3.43 (q, 2H), 2.78 (m, 1H), 2.33 (m, 2H), 2.02 (m, 5H), 1.99 (m, 4H), 1.60 (m, 1H), 1.26 (m, 1H)

#### Example 16:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-nitrophenyl)piperidin-3-one

In accordance with the same procedure as Example 1, except that 4-fluoronitrobenzene was used instead of 1-fluorophenyl methyl sulfone in Step 3 thereof, 190 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.13(m, 1H), 7.68(m, 1H), 7.54(m, 2H), 7.47(m, 2H), 7.45 (m, 2H), 6.75(s, 1H), 4.78 (m, 1H), 4.24 (d, 1H), 4.03(d, 1H), 3.89(t, 1H) 2.94 (m,1H), 2.27 (m, 3H), 1.98 (m, 5H), 1.76(m, 5H), 1.72(m, 3H), 1.52 (m, 1H)

20

#### Example 17:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(3-methyl-4-nitrophenyl)piperidin-3-one

25

In accordance with the same procedure as in Example 1, except that 4-fluoro-2-methyl-nitrobenzene was used instead of 1-fluorophenyl methyl sulfone in Step 3 thereof, 170 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.69 (d, 1H), 7.55 (d, 1H), 7.53 (m, 3H), 7.33 (m, 1H), 7.14 (m, 1H), 6.73 (m, 3H), 4.72 (m, 1H), 3.59 (d, 1H), 3.18 (s, 3H), 3.15 (m, 2H), 2.75 (m, 1H), 2.64 (s, 3H), 2.34 (m, 2H), 2.16 (m, 2H), 1.98(m, 1H), 1.79 (m, 3H), 1.72 (m, 2H), 1.52 (m, 1H)

30

Example 18:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(3-fluoro-2-formylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 1, except that 2,6-difluorobenzaldehyde was used instead of 1-fluorophenyl methyl sulfone in Step 3 thereof, 210 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 10.33 (s, 1H), 7.94 (d, 1H), 7.61(m, 1H), 7.35 (m, 1H), 7.23 (m, 2H), 7.04 (m, 1H), 6.99 (m, 1H), 4.78 (m, 1H), 4.12 (m, 1H), 3.76 (m, 1H), 3.25 (m, 1H), 3.07 (m, 1H), 2.94 (m, 1H), 2.36 (m, 1H), 2.22 (m, 1H), 2.01 (m, 4H), 1.63 (m, 3H), 1.45 (m, 1H), 1.24 (m, 1H)

Example 19:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(3-chloro-2-formylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 1, except that 2-chloro-6-fluorobenzaldehyde was used instead of 1-fluorophenyl methyl sulfone in Step 3 thereof, 200 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 10.31 (s, 1H), 7.66 (d, 1H), 7.53 (m, 1H), 7.34 (m, 1H), 7.26 (m, 2H), 6.95 (m, 1H), 6.93 (m, 1H), 4.69 (m, 1H), 4.11 (m, 1H), 3.39 (m, 2H), 3.25 (m, 1H), 2.23 (m, 1H), 2.32 (m, 1H), 2.05 (m, 1H), 2.01 (m, 4H), 1.72 (m, 3H), 1.53 (m, 1H), 1.26 (m, 1H)

Example 20:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-formyl-4-nitrophenyl)piperidin-3-one

In accordance with the same procedure as in Example 1, except that 2-fluoro-5-nitrobenzaldehyde was used instead of 1-fluorophenyl methyl sulfone in Step 3 thereof, 180 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.63 (s, 1H), 8.23 (d, 1H), 7.69 (m, 2H),  
5 7.53 (m, 2H), 7.47 (m, 3H), 6.84 (s, 1H), 4.84 (m, 1H), 4.51 (m, 1H), 4.32 (m,  
1H), 2.35 (m, 1H), 3.32 (m, 3H), 2.06 (m, 3H), 1.78 (m, 4H), 1.55 (m, 1H), 1.25  
(m, 1H)

#### Example 21:

10 Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-for  
myl-6-fluorophenyl)piperidin-3-one

In accordance with the same procedure as in Example 1, except that  
15 2,3-difluorobenzaldehyde was used instead of 1-fluorophenyl methyl sulfone in  
Step 3 thereof, 110 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 10.42 (s, 1H), 7.86 (d, 1H), 7.65 (m, 2H),  
7.61 (m, 1H), 7.30 (m, 2H), 7.02 (m, 1H), 6.42 (s, 1H), 4.37 (m, 1H), 3.56 (m,  
1H), 3.46 (m, 2H), 3.03 (m, 1H), 2.45 (m, 1H), 2.21 (m, 1H), 2.02 (m, 2H), 1.94  
20 (m, 1H), 1.76 (m, 4H), 1.64 (m, 3H), 1.28 (m, 2H)

#### Example 22:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-flu  
25 oro-3-formylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 1, except that  
2,3-difluorobenzaldehyde was used instead of 1-fluorophenyl methyl sulfone in  
Step 3 thereof, 100 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 10.41(s, 1H), 7.80 (d, 1H), 7.59 (m, 1H),  
30 7.34 (m, 1H), 7.26 (m, 2H), 7.17 (m, 1H), 6.81 (m, 1H), 4.15 (m, 1H), 4.13 (m,  
1H), 3.95 (m, 1H), 3.36 (m, 2H), 3.15 (m, 1H), 3.00 (m, 1H), 2.29 (m, 3H), 2.01

(m, 5H), 1.79 (m, 4H), 1.54 (m, 3H), 1.28 (m, 2H)

Example 23:

Preparation of

5 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-isopropylsulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 1, except that 1-fluorophenyl isopropyl sulfone was used instead of 1-fluorophenyl methyl sulfone in Step 3 thereof, 170 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.05 (d, 1H), 7.69 (d, 1H), 7.63 (t, 1H), 7.54 (d, 1H), 7.43 (m, 3H), 7.29 (m, 3H), 6.73 (s, 1H), 4.70 (m, 1H), 3.77 (m, 1H), 3.76 (d, 1H), 3.57 (d, 1H), 3.54 (m, 1H), 3.15 (t, 1H), 2.73 (m, 1H), 2.31 (t, 2H), 2.05 (t, 2H), 1.90 (m, 1H), 1.74 (m, 3H), 1.56 (m, 2H), 1.40 (m, 1H) 1.33 (d, 3H), 1.06 (d, 3H)

Example 24:

Preparation of

20 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-trifluoromethylphenyl)piperidin-3-one

Step 1.

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-trifluoromethylphenyl)piperidin-3-ol

25 To the suspension of 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-3-piperidinol hydrochloride (300 mg) obtained in Step 2 of Example 1 in 5 ml of toluene, were added 1-bromo-4-fluoro-2-trifluoromethylbenzene (0.198 ml), cesium carbonate (1.42 g), (S)-2,2-bis(diphenylphosphino)-1,1-binaphthyl (BINAP; 45.4 mg) and tris(dibenzylideneacetone)dipalladium (1.42 g). The reaction mixture was stirred for about 4 hours at 100 °C under nitrogen atmosphere, was diluted

with 30ml of ethyl acetate, filtered, and concentrated. The resulting residue was purified with column chromatography to give 179 mg of the titled compound.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.80-7.00 (m, 6H), 7.70 (m, 1H), 4.80 (m, 1H), 3.80-3.60 (m, 3H), 3.30 (m, 1H), 3.00 (m, 1H), 2.69 (m, 1H), 2.33 (d, 2H), 2.01 (t, 2H), 1.80-1.68 (m, 2H), 1.43 (t, 2H), 1.35 (d, 2H)

#### Step 2.

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-trifluoromethylphenyl)piperidin-3-one

To the solution of 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-trifluoromethylphenyl)piperidin-3-ol (0.15 g) obtained in Step 1 in 3 ml of dimethyl sulfoxide (DMSO), was added 0.16 ml of triethylamine. The reaction mixture was cooled to 0°C. Pyridine-SO<sub>3</sub> complex (0.15 g) was added slowly to the reaction mixture, which was then stirred for about 4 hours at room temperature, diluted with 20 ml of ethyl acetate, washed with brine, and dried and concentrated. The resulting residue was purified with column chromatography to give 84 mg of the titled compound as white solid.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.69 (d, 1H), 7.56-7.27 (m, 8H), 6.75 (s, 1H), 4.68 (m, 1H), 8.95 (d, 1H), 3.71 (m, 2H), 3.39 (t, 1H), 2.78 (m, 1H), 2.31 (d, 2H), 2.03 (m, 2H), 1.87 (m, 1H), 1.75 (m, 3H), 1.53 (m, 2H), 1.39 (m, 1H)

#### Example 25:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-cyanophenyl)piperidin-3-one

In accordance with the same procedure as in Example 24, except that 2-bromobenzonitrile was used instead of 1-bromo-4-fluoro-2-trifluoromethylbenzene in Step 1 thereof, 160 mg of the

titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.69 (d, 1H), 7.57 (m, 2H), 7.49 (m, 2H),  
7.31 (m, 1H), 7.06 (t, 1H), 6.97 (d, 1H), 6.73 (s, 1H), 4.71 (m, 1H), 3.86 (q, 3H),  
3.41 (t, 1H), 2.78 (m, 1H), 2.31 (broad, 2H), 2.00 (m, 3H), 1.75 (m, 2H), 1.52 (m,  
5 2H), 1.38 (m, 1H)

Example 26:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(4-  
10 morpholino)phenyl]piperidin-3-one

In accordance with the same procedure as in Example 24, except that  
2-(4-morpholino)bromobenzene was used instead of  
1-bromo-4-fluoro-2-trifluoromethylbenzene in Step 1 thereof, 110 mg of the  
15 titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.75 (d, 1H), 7.55 (d, 1H), 7.43 (m, 3H),  
7.34 (t, 1H), 7.12 (m, 2H), 6.93 (m, 2H), 6.74 (s, 1H), 4.65 (m, 1H), 4.21 (d, 1H),  
3.84 (m, 4H), 3.50 (d, 1H), 3.13 (m, 1H), 3.08 (m, 4H), 2.81 (m, 1H), 2.32 (d,  
20 2H), 2.12 (t, 2H), 1.75 (m, 4H), 1.52 (m, 2H), 1.32 (m, 1H), 1.25 (m, 1H)

Example 27:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-et  
25 hoxyphenyl)piperidin-3-one

In accordance with the same procedure as in Example 24, except that  
2-bromoethoxybenzene was used instead of  
1-bromo-4-fluoro-2-trifluoromethylbenzene in Step 1 thereof, 95 mg of the titled  
compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.75 (d, 1H), 7.65 (d, 1H), 7.52 (m, 3H),  
30 7.39 (t, 1H), 7.04 (m, 1H), 6.85 (m, 2H), 6.74 (s, 1H), 4.64 (m, 1H), 4.05 (q, 2H),  
3.97 (d, 1H), 3.65 (m, 1H), 3.58 (d, 1H), 3.21 (m, 1H), 2.72 (m, 1H), 2.29 (m,

2H), 2.07 (t, 2H), 1.81 (m, 1H), 1.65 (m, 3H), 1.52 (t, 2H), 1.35 (t, 3H), 1.32 (m, 1H)

Example 28:

5 Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-fluoro-4-trifluoromethylphenyl)piperidin-3-one

10 In accordance with the same procedure as in Example 24, except that 1-bromo-2-fluoro-4-trifluoromethylbenzene was used instead of 1-bromo-4-fluoro-2-trifluoromethylbenzene in Step 1 thereof, 95 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.69 (d, 1H), 7.55 (d, 1H), 7.47 (m, 3H), 7.33 (m, 2H), 7.26 (s, 1H), 6.96 (t, 1H), 6.73 (s, 1H), 4.71 (m, 1H), 8.95 (d, 1H), 15 3.71 (m, 2H), 3.39 (t, 1H), 2.78 (m, 1H), 2.31 (d, 2H), 2.03 (m, 2H), 1.87 (m, 1H), 1.75 (m, 3H), 1.5 (m, 1H)

Example 29:

Preparation of

20 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-chloro-4-trifluoromethylphenyl)piperidin-3-one

25 In accordance with the same procedure as in Example 24, except that 1-bromo-2-chloro-4-trifluoromethylbenzene was used instead of 1-bromo-4-fluoro-2-trifluoromethylbenzene in Step 1 thereof, 120 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.69 (d, 1H), 7.56-7.27 (m, 8H), 6.75 (s, 1H), 4.68 (m, 1H), 8.95 (d, 1H), 3.71 (m, 2H), 3.39 (t, 1H), 2.78 (m, 1H), 2.31 (d, 2H), 2.03 (m, 2H), 1.87 (m, 1H), 1.75 (m, 3H), 1.53 (m, 2H), 1.39 (m, 1H)

30

Example 30:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2,4-dichlorophenyl)piperidin-3-one

In accordance with the same procedure as in Example 24, except that  
5 1-bromo-2,4-dichlorobenzene was used instead of  
1-bromo-4-fluoro-2-trifluoromethylbenzene in Step 1 thereof, 100 mg of the  
titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.69 (d, 1H), 7.56 (d, 1H), 7.49-7.45 (m,  
3H), 7.35-7.31 (m, 2H), 7.19 (d, 1H), 6.93 (d, 1H), 6.74 (s, 1H), 4.69 (m, 1H),  
10 3.74 (d, 1H), 3.71 (d, 1H), 3.69 (d, 1H), 3.19 (t, H), 2.88 (m, 1H), 2.31 (m, 2H),  
2.02 (m, 2H), 1.91 (m, 1H), 1.75 (m, 2H), 1.54 (t, 2H), 1.40 (m, 1H)

Example 31:

Preparation of

15 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-ch  
loro-2-fluorophenyl)piperidin-3-one

In accordance with the same procedure as in Example 24, except that  
1-bromo-4-chloro-2-fluorobenzene was used instead of  
20 1-bromo-4-fluoro-2-trifluoromethylbenzene in Step 1 thereof, 80 mg of the titled  
compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.69 (d, 1H), 7.54 (d, 1H), 7.47-7.45 (m,  
3H), 7.31 (t, 1H), 7.05 (m, 2H), 6.84 (t, 1H), 6.72 (s, 1H), 4.68 (m, 1H), 3.83 (d,  
1H), 3.79 (d, 1H), 3.52 (d, 1H), 3.29 (t, H), 2.78 (m, 1H), 2.31 (m, 2H), 2.02 (m,  
25 2H), 1.91 (m, 1H), 1.75 (m, 2H), 1.54 (t, 2H), 1.40 (m, 1H)

Example 32:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-m  
30 ethoxy-5-cyanophenyl)piperidin-3-one

In accordance with the same procedure as in Example 24, except that



3-bromo-4-methoxybenzonitrile was used instead of 1-bromo-4-fluoro-2-trifluoromethylbenzene in Step 1 thereof, 50 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.69 (d, 1H), 7.54 (d, 1H), 7.47-7.45 (m, 3H), 7.31 (t, 1H), 7.05 (m, 2H), 6.84 (t, 1H), 6.72 (s, 1H), 4.69 (m, 1H), 3.89 (s, 3H), 3.88 (d, 1H), 3.84 (d, 1H), 3.56 (d, 1H), 3.23 (t, H), 2.78 (m, 1H), 2.31 (m, 2H), 2.02 (m, 2H), 1.91 (m, 1H), 1.75 (m, 2H), 1.54 (t, 2H), 1.40 (m, 1H)

Example 33:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-cyano-2-fluorophenyl)piperidin-3-one

In accordance with the same procedure as in Example 24, except that 4-bromo-3-fluorobenzonitrile was used instead of 1-bromo-4-fluoro-2-trifluoromethylbenzene in Step 1 thereof, 110 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.69 (d, 1H), 7.56 (d, 1H), 7.49-7.45 (m, 3H), 7.35-7.31 (m, 2H), 7.19 (d, 1H), 6.93 (d, 1H), 6.74 (s, 1H), 4.69 (m, 1H), 3.74 (d, 1H), 3.71 (d, 1H), 3.69 (d, 1H), 3.19 (t, H), 2.88 (m, 1H), 2.31 (m, 2H), 2.02 (m, 2H), 1.91 (m, 1H), 1.75 (m, 2H), 1.54 (t, 2H), 1.40 (m, 1H)

Example 34:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(2-furyl)phenyl]piperidin-3-one

In accordance with the same procedure as in Example 24, except that 1-bromo-2-(2-furyl)benzene was used instead of 1-bromo-4-fluoro-2-trifluoromethylbenzene in Step 1 thereof, 75 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.66 (d, 1H), 7.55-7.44 (m, 5H), 7.42-7.31

(m, 4H), 7.29-7.11 (m, 2H), 7.03 (d, 1H), 6.96 (t, 1H), 6.74 (m, 2H), 4.70 (m, 1H), 3.74 (d, 1H), 3.71 (d, 1H), 3.69 (d, 1H), 3.19 (t, 1H), 2.88 (m, 1H), 2.31 (m, 2H), 2.02 (m, 2H), 1.40 (m, 1H)

5 Example 35:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-bromophenyl)piperidin-3-one

10 In accordance with the same procedure as in Example 24, except that 1,2-dibromobenzene was used instead of 1-bromo-4-fluoro-2-trifluoromethylbenzene in Step 1 thereof, 85 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.75 (d, 1H), 7.55 (m, 4H), 7.35 (m, 3H),  
15 7.15 (d, 1H), 6.88 (m, 1H), 6.77 (s, 1H) 4.72 (m, 1H), 3.88 (d, 1H), 3.74 (d, 1H), 3.61 (m, 1H), 2.75 (m, 1H), 3.33 (d, 2H), 2.02 (t, 2H), 1.88 (m, 1H), 1.76 (m, 3H), 1.55 (m, 1H), 1.44 (m, 1H)

Example 36:

20 Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methylthiophenyl)piperidin-3-one

In accordance with the same procedure as in Example 24, except that  
25 2-bromothioanisole was used instead of 1-bromo-4-fluoro-2-trifluoromethylbenzene in Step 1 thereof, 65 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.75 (d, 1H), 7.68 (m, 4H), 7.35 (m, 2H),  
7.15 (m, 4H), 6.73 (s, 1H), 4.65 (m, 1H), 3.75 (d, 1H), 3.66 (d, 1H), 3.40 (m, 1H),  
30 3.17 (t, 1H) 2.72 (m, 1H), 2.39 (s, 3H), 2.33 (m, 2H), 2.09 (t, 2H), 1.87 (m, 1H), 1.72 (m, 3H), 1.55 (m, 1H) 1.44 (m, 1H)

Example 37:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(tert-butoxycarbonyl)phenyl]piperidin-3-one

5

In accordance with the same procedure as in Example 24, except that tert-butyl 2-bromobenzoate was used instead of 1-bromo-4-fluoro-2-trifluoromethylbenzene in Step 1 thereof, 65 mg of the titled compound was prepared.

10  $^1\text{H-NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  7.64 (m, 2H), 7.51 (d, 1H), 7.47 (m, 2H), 7.44 (m, 3H), 6.97 (d, 1H), 6.89 (m, 1H), 6.72 (s, 1H), 4.63 (m, 1H), 3.80 (d, 1H), 3.77 (d, 1H), 3.63 (m, 1H), 3.29 (m, 1H), 2.61 (m, 1H), 2.29 (d, 2H), 1.97 (m, 3H), 1.76 (m, 3H), 1.58 (s, 9H), 1.50 (m, 2H), 1.38 (m, 1H)

15 Example 38:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-ethoxycarbonylphenyl)piperidin-3-one

20 In accordance with the same procedure as in Example 24, except that ethyl 2-bromobenzoate was used instead of 1-bromo-4-fluoro-2-trifluoromethylbenzene in Step 1 thereof, 80 mg of the titled compound was prepared.

25  $^1\text{H-NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  7.64 (m, 2H), 7.51 (d, 1H), 7.47 (m, 2H), 7.44 (m, 3H), 6.97 (d, 1H), 6.89 (m, 1H), 6.72 (s, 1H), 4.63 (m, 1H), 4.30 (q, 2H), 3.80 (d, 1H), 3.77 (d, 1H), 3.63 (m, 1H), 3.29 (m, 1H), 2.61 (m, 1H), 2.29 (d, 2H), 1.97 (m, 3H), 1.76 (m, 3H), 1.50 (m, 2H), 1.38 (m, 1H), 1.20 (t, 3H)

Example 39:

30 Preparation of  
4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methoxyphenyl)piperidin-3-one

In accordance with the same procedure as in Example 24, except that 2-bromoanisole was used instead of 1-bromo-4-fluoro-2-trifluoromethylbenzene in Step 1 thereof, 60 mg of the titled compound was prepared.

5       <sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.75 (d, 1H), 7.67 (d, 1H), 7.51 (m, 3H), 7.34 (m, 1H), 7.07 (m, 1H), 6.91 (m, 3H), 6.74 (s, 1H), 4.66 (m, 1H), 3.92 (d, 1H), 3.84 (s, 3H), 3.58 (m, 2H), 3.22 (t, 1H), 2.75 (m, 1H) 2.32 (m, 2H), 2.05 (m, 2H), 1.85 (m, 1H), 1.75 (m, 3H), 1.52 (m, 2H), 1.35 (m, 1H)

10       Example 40:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-methylphenyl)piperidin-3-one

15       In accordance with the same procedure as in Example 24, except that 2-bromo-5-fluorotoluene was used instead of 1-bromo-4-fluoro-2-trifluoromethylbenzene in Step 1 thereof, 70 mg of the titled compound was prepared.

20       <sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.68 (d, 1H), 7.48 (m, 4H), 7.33 (m, 1H), 6.98 (m, 1H), 6.85 (m, 2H), 6.75 (s, 1H), 4.65 (m, 1H), 3.51 (s, 2H), 3.13 (d, 2H), 2.74 (m, 1H), 2.33 (m, 2H), 2.26 (s, 3H), 2.01 (m, 3H), 1.75 (m, 3H), 1.52 (t, 2H), 1.32 (m, 1H)

Example 41:

25       Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-isopropoxyphenyl)piperidin-3-one

30       In accordance with the same procedure as in Example 24, except that 1-bromo-2-isopropoxybenzene was used instead of 1-bromo-4-fluoro-2-trifluoromethylbenzene in Step 1 thereof, 55 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.65 (d, 1H), 7.59-7.37(m, 4H), 7.27 (t, 1H), 6.97 (m, 1H), 6.85 (m, 3H), 6.72 (s, 1H), 4.65 (m, 1H), 4.55 (m, 1H), 3.89 (d, 1H), 3.58 (m, 2H), 3.21 (t, 1H), 2.68 (m, 1H), 2.31 (m, 2H), 2.01 (m, 2H), 1.72 (m, 4H), 1.49 (m, 2H), 1.41 (m, 1H), 1.32 (dd, 6H)

5

Example 42:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-ethylthiophenyl)piperidin-3-one

10

In accordance with the same procedure as in Example 24, except that 1-bromo-2-ethylthiobenzene was used instead of 1-bromo-4-fluoro-2-trifluoromethylbenzene in Step 1 thereof, 40 mg of the titled compound was prepared.

15

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.75 (d, 1H), 7.59-7.41 (m, 4H), 7.33 (m, 1H), 7.25 (m, 3H), 7.05 (d, 1H), 6.73 (s, 1H), 4.65 (m, 1H), 3.69 (d, 1H), 3.55 (d, 1H), 3.45 (d, 1H), 3.15 (t, 1H), 2.89 (q, 2H), 2.71 (m, 1H), 2.31 (d, 2H), 2.05 (t, 2H), 1.89 (m, 1H), 1.75 (m, 3H), 1.54 (m, 2H), 1.36 (m, 1H) 1.34 (t, 3H)

20 Example 43:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(3,3-dimethylazetidin-2-one-1-yl)phenyl]piperidin-3-one

25 Step 1.

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-nitrophenyl)piperidin-3-ol

To the solution of  
30 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-3-piperidinol hydrochloride (6.0 g) obtained in Step 2 of Example 1 in 50 ml of ethanol, were added 3.7 ml of 2-fluoronitrobenzene and 4.8 ml of triethylamine. The

reaction mixture was refluxed overnight and concentrated. The resulting residue was filtered and dried to give 7.2 g of the titled compound.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.70 (d, 1H), 7.68-7.46 (m, 4H), 7.33 (t, 1H), 7.21-7.15 (m, 2H), 6.99 (d, 1H), 6.92 (d, 1H), 6.84 (d, 1H), 6.82 (s, 1H),  
5 4.09 (broad, 1H), 3.81 (m, 2H), 3.67 (m, 1H), 3.31 (d, 1H), 3.15 (d, 1H), 2.66 (m, 2H), 2.28 (t, 2H), 2.02 (m, 2H), 1.76-1.55 (m, 4H), 1.48 (t, 2H), 1.28 (m, 1H)

#### Step 2.

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-aminophenyl)piperidin-3-ol  
10

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-nitrophenyl)piperidin-3-ol (7.0 g) obtained in Step 1 was dissolved in 30 ml of methanol. The solution was stirred with 10%-palladium catalyst overnight  
15 under 30 psi hydrogen atmosphere, and then filtered and concentrated to give 6.5 g of titled compound.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.70 (d, 1H), 7.68-7.52 (m, 2H), 7.46 (m, 1H), 7.33 (m, 1H), 7.15 (d, 1H), 6.99 (d, 1H), 6.92 (d, 1H), 6.72 (s, 1H), 6.70 (m, 3H), 4.09 (broad, 1H), 3.81 (m, 2H), 3.67 (m, 1H), 3.31 (d, 1H), 3.15 (d, 1H),  
20 2.66 (m, 2H), 2.28 (t, 2H), 2.02 (m, 2H), 1.76-1.55 (m, 4H), 1.48 (t, 2H), 1.28 (m, 1H)

#### Step 3.

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(3-chloro-2,2-dimethylbutyryl)aminophenyl]piperidin-3-ol  
25

The solution of 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-aminophenyl)piperidin-3-ol (4.0 g) obtained in Step 2 in 30 ml of dichloromethane was cooled to 0°C. 2.2 ml of triethylamine and 2.1 ml of  
30 3-chloropivaloyl chloride were added dropwise to the reaction mixture, which was then stirred for about 2 hours at room temperature. 20 ml of iced water was added to the reaction mixture. The resulting organic layer was dried and

concentrated on sodium sulfate. The resulting residue was crystallized with ethyl acetate to give 4.1 g of the titled compound.

Step 4.

5 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(3, 3-dimethylazetidin-2-one-1-yl)phenyl]piperidin-3-ol

The solution of 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(3-  
10 chloro-2,2-dimethylbutyryl)aminophenyl]piperidin-3-ol (4.1 g) obtained in Step 3 in 50 ml of dimethylformamide was cooled to 0 °C. Sodium hydride (NaH, 60%; 640 mg) was added to the reaction mixture, which was then stirred overnight at 50 °C, diluted with ethyl acetate, washed with a brine, and then dried and concentrated on sodium sulfate. The resulting residue was purified with  
15 column chromatography to give 2.1 g of the titled compound.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.72 (t, 2H), 7.54 (m, 3H), 7.33 (t, 1H), 7.07 (m, 4H), 6.77 (s, 1H), 4.22 (d, 1H), 3.81 (m, 1H), 3.73 (m, 2H), 3.66 (d, 1H), 3.56 (d, 1H), 3.37 (d, 1H), 3.11 (d, 1H), 2.71 (m, 2H), 2.28 (m, 2H), 2.06 (m, 2H), 1.78 (m, 4H), 1.38 (m, 2H), 1.23 (d, 6H)

20

Step 5.

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(3, 3-dimethylazetidin-2-one-1-yl)phenyl]piperidin-3-one

25 To the solution of 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(3, 3-dimethylazetidin-2-one-1-yl)phenyl]piperidin-3-ol (0.14 g) obtained in Step 4 in 3 ml of dimethylsulfoxide (DMSO), was added 0.16 ml of triethylamine. The reaction mixture was cooled to 0 °C. Pyridine- SO<sub>3</sub> complex (0.15 g) was  
30 added slowly to the reaction mixture, which was stirred for about 10 hours at room temperature. The reaction mixture was diluted with 20 ml of ethyl

acetate, washed with brine. The resulting organic layer was dried and concentrated on sodium sulfate. The resulting residue was filtered and dried to give 120 mg of the titled compound as white color.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.87 (d, 1H), 7.70 (d, 1H), 7.56 (d, 1H),  
5 7.50-7.46 (m, 3H), 7.33 (d, 1H), 7.11 (t, 3H), 6.74 (s, 1H), 4.67 (t, 1H), 3.62 (t, 2H), 3.46 (t, 2H), 2.69 (m, 1H), 2.33 (d, 2H), 2.01 (t, 2H), 1.80-1.68 (m, 2H), 1.43 (t, 2H), 1.35 (d, 6H)

#### Example 44:

10 Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(p  
yrrolidin-2-one-1-yl)phenyl]piperidin-3-one

In accordance with the same procedure as in Example 43, except that  
15 4-chlorobutyl chloride was used instead of 3-chloropivaloyl chloride in Step 3 thereof, 105 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.68 (d, 1H), 7.54 (d, 1H), 7.50-7.36 (m,  
3H), 7.32 (t, 1H), 7.25-7.17 (m, 4H), 7.11 (t, 1H), 7.08 (d, 1H), 6.72 (s, 1H), 4.65  
(m, 1H), 3.73 (q, 1H), 3.66 (m, 1H), 3.63 (d, 1H), 3.55 (d, 1H), 3.42 (d, 1H),  
20 3.28 (t, 1H), 2.78 (m, 1H), 2.57 (m, 2H), 2.30 (d, 2H), 2.13 (m, 2H), 2.01 (t, 2H), 1.76 (m, 4H), 1.48 (t, 2H), 1.24 (m, 1H)

Mass (M+H) = 543

#### Example 45:

25 Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(pi  
peridin-2-one-1-yl)phenyl]piperidin-3-one

In accordance with the same procedure as in Example 43, except that  
30 5-chlorovaleryl chloride was used instead of 3-chloropivaloyl chloride in Step 3 thereof, 90 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.68 (d, 1H), 7.54 (d, 1H), 7.50-7.36 (m,



3H), 7.32 (t, 1H), 7.25-7.17 (m, 4H), 7.11 (t, 1H), 7.08 (d, 1H), 6.72 (s, 1H), 4.65 (m, 1H), 3.73 (q, 1H), 3.66 (m, 1H), 3.63 (d, 1H), 3.55 (d, 1H), 3.42 (d, 1H), 3.28 (t, 1H), 2.78 (m, 1H), 2.57 (m, 2H), 2.30 (d, 2H), 2.13 (m, 2H), 2.01 (t, 2H), 1.76 (m, 4H), 1.48 (t, 2H), 1.10-1.30 (m, 3H)

5

Example 46:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(2-oxazolidinon-1-yl)phenyl]piperidin-3-one

10

In accordance with the same procedure as in Example 43, except that 2-chloroethyl chloroformate was used instead of 3-chloropivaloyl chloride in Step 3 thereof, 75 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.70 (d, 1H), 7.68 (m, 4H), 7.33 (m, 2H),  
15 7.26 (m, 1H), 7.16 (m, 1H), 6.72 (s, 1H), 4.70 (m, 1H), 4.50 (m, 2H), 4.01 (q, 1H), 3.92 (q, 1H), 3.72 (d, 1H), 3.69 (d, 1H), 3.36 (m, 1H), 3.15 (m, 1H), 2.74 (m, 1H), 2.31 (m, 2H), 2.01 (m, 2H), 1.78 (m, 4H)

Example 47:

20 Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(2-imidazolidinon-1-yl)phenyl]piperidin-3-one

In accordance with the same procedure as in Example 43, except that  
25 2-chloroethyl isocyanate was used instead of 3-chloropivaloyl chloride in Step 3 thereof, 80 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.69 (d, 1H), 7.60 (d, 1H), 7.50 (m, 2H),  
7.34 (m, 2H), 7.31 (m, 1H), 7.03 (m, 1H), 7.01 (m, 1H), 6.74 (s, 1H), 4.93 (s, 1H), 4.66 (m, 1H), 3.86 (m, 2H), 3.50 (m, 4H), 3.15 (m, 1H), 2.76 (m, 1H), 2.33  
30 (m, 2H), 2.00 (m, 2H), 1.77 (m, 4H), 1.52 (m, 2H), 1.20 (m, 1H)

Example 48:

## Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-[N-(ethoxycarbonyl)amino]phenyl]piperidin-3-one

- 5      Step 1.  
4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-[N-(ethoxycarbonyl)amino]phenyl]piperidin-3-ol

The solution of 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexane  
10 carbonyl]amino]-1-(2-aminophenyl)piperidin-3-one (0.9 g) obtained in Step 2 of  
Example 43 in 20 ml of dichloromethane was cooled to 0°C. 0.54 ml of  
triethylamine and 0.37 ml of ethyl chloroformate were added to the reaction  
mixture, which was then stirred for about 3 hours at room temperature. 20 ml  
of ice water was added to the reaction mixture. The resulting organic layer  
15 was dried and concentrated on sodium sulfate. The resulting residue was  
purified with column chromatography to give 600 mg of the titled compound.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.12 (d, 1H), 7.75 (d, 1H), 7.57 (m, 2H),  
7.50 (m, 1H), 7.13 (m, 1H), 7.11 (m, 2H), 7.02 (m, 1H), 6.86 (s, 1H), 4.25 (s,  
1H), 4.20 (q, 2H), 3.85 (m, 1H), 3.62 (m, 1H), 3.39 (m, 1H), 3.04 (m, 1H), 2.73  
20 (m, 2H), 2.24 (m, 2H), 2.09 (m, 2H), 1.96 (m, 4H), 1.73 (m, 3H), 1.66 (m, 2H),  
1.25 (m, 1H), 1.23 (t, 3H)

- Step 2.  
4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-[N-  
25 -(ethoxycarbonyl)amino]phenyl]piperidin-3-one

To the solution of 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]  
cyclohexanecarbonyl]amino]-1-[2-[N-(ethoxycarbonyl)amino]phenyl]piperidin-3-  
ol (600mg, 1.1 mmol) in 10 ml of dimethylsulfoxide (DMSO), was added 0.46  
30 ml of triethylamine (3.3 mmol). The reaction mixture was cooled to 0°C.  
Pyridine-SO<sub>3</sub> complex (0.51 g, 3.3 mmol) was added to the reaction mixture,  
which was then stirred for about 2 hours at room temperature. The reaction

mixture was diluted with 20 ml of ethyl acetate, washed with brine. The resulting organic layer was dried and concentrated on sodium sulfate. The resulting residue was filtered and dried to give 320 mg of the titled compound as white color.

5       <sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.12 (d, 1H), 7.57 (d, 1H), 7.55 (m, 3H), 7.54 (m, 1H), 7.50 (m, 1H), 7.12 (m, 2H), 6.86 (s, 1H), 4.65 (m, 1H), 4.20 (q, 2H), 3.54 (d, 1H), 3.33 (d, 1H), 3.12 (m, 2H), 2.76 (m, 1H), 2.25 (m, 2H), 2.03 (m, 2H), 1.89 (m, 1H), 1.68 (m, 3H), 1.45 (m, 2H), 1.30 (m, 1H), 1.26 (t, 3H)

10    Example 49:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-[N-(methoxycarbonyl)amino]phenyl]piperidin-3-one

15       In accordance with the same procedure as in Example 48, except that methyl chloroformate was used instead of ethyl chloroformate in Step 1 thereof, 305 mg of the titled compound was prepared.

20       <sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.11 (d, 1H), 7.69(d, 1H), 7.54 (m, 3H), 7.52 (m, 1H), 7.33 (m, 1H), 7.31 (m, 2H), 7.03 (m, 1H), 6.76 (s, 1H), 4.66 (m, 1H), 3.75 (s, 3H), 3.58 (m, 5H), 1.76 (m, 2H), 1.32 (m, 1H)

Example 50:

Preparation of 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-[N-(4-chlorobutyryl)amino]phenyl]piperidin-3-one

25

In accordance with the same procedure as in Example 48, except that 4-chlorobutyryl chloride was used instead of ethyl chloroformate in Step 1 thereof, 290 mg of the titled compound was prepared.

30       <sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.69 (d, 1H), 7.56 (d, 1H), 7.49-7.45 (m, 3H), 7.35-7.31 (m, 2H), 7.19 (d, 1H), 7.10 (d, 1H), 6.93 (d, 1H), 6.73 (s, 1H), 4.69 (m, 1H), 3.66 (m, 2H), 3.58 (d, 1H), 3.49 (d, 1H), 3.23 (d, 1H), 3.09 (t, H), 2.55 (m, 2H), 2.48 (t, 2H), 2.31 (m, 2H), 2.12 (m, 2H), 2.02 (m, 2H), 1.91 (m,

1H), 1.75 (m, 2H), 1.54 (t, 2H), 1.40 (m, 1H)

Example 51:

Preparation of

5 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-[N-(isobutyryl)amino]phenyl]piperidin-3-one

In accordance with the same procedure as in Example 48, except that isobutyryl chloride was used instead of ethyl chloroformate in Step 1 thereof,  
10 330 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.20 (d, 1H), 7.66 (d, 1H), 7.46-7.43 (m, 3H), 7.32-7.28 (m, 2H), 7.11-7.05 (m, 1H), 6.77 (d, 1H), 4.75 (m, 1H), 3.80-3.76 (m, 1H), 3.74 (m, 1H), 3.67-3.63 (m, 2H), 3.30-3.25 (m, 1H), 3.17-3.14 (m, 1H), 2.87-2.65 (m, 2H), 2.60-2.54 (m, 2H), 2.19-2.05 (m, 2H), 1.77-1.71 (m, 2H),  
15 1.57-1.41 (m, 1H), 1.42-1.30 (m, 1H), 1.20-1.12 (m, 7H)

Example 52:

Preparation of

20 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-[N-(methoxyacetyl)amino]phenyl]piperidin-3-one

In accordance with the same procedure as in Example 48, except that methoxyacetyl chloride was used instead of ethyl chloroformate in Step 1 thereof, 250 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.68 (d, 1H), 7.55-7.43 (m, 3H), 7.35-7.30 (m, 2H), 7.14-7.02 (m, 1H), 6.80(m, 1H), 4.81-4.75 (m, 1H), 3.80-3.76 (m, 1H), 3.67-3.63 (m, 2H), 3.48 (s, 3H), 3.45-3.25 (m, 3H), 3.17-3.14 (m, 1H), 2.78-2.72 (m, 1H), 2.55-2.50 (m, 2H), 2.36-2.30 (m, 1H), 2.23-2.12 (m, 2H), 1.77-1.71 (m, 2H), 1.57-1.41 (m, 1H), 1.30-1.25 (m, 1H)

Example 53:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-[N-(cyclopropanecarbonyl)amino]phenyl] piperidin-3-one

In accordance with the same procedure as in Example 48, except that  
5 cyclopropanecarbonyl chloride was used instead of ethyl chloroformate in Step 1 thereof, 150 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.70 (d, 1H), 7.50-7.40 (m, 3H),  
7.35-7.05 (m, 3H), 6.85 (m, 1H), 4.78-4.75 (m, 1H), 3.80-3.76 (m, 1H),  
3.76-3.70 (m, 1H), 3.65-3.60 (m, 2H), 3.30-3.25 (m, 1H), 3.17-3.14 (m, 1H),  
10 2.78-2.72 (m, 1H), 2.57-2.51 (m, 2H), 2.40-2.28 (m, 2H), 2.20-2.00 (m, 3H),  
1.98-1.62 (m, 4H), 1.57-1.41 (m, 3H), 1.30-1.20 (m, 1H)

Example 54:

Preparation of

15 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-[N-(3-phenylpropionyl)amino]phenyl]piperidin-3-one

In accordance with the same procedure as in Example 48, except that  
20 3-phenylpropionyl chloride was used instead of ethyl chloroformate in Step 1 thereof, 220 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.34 (d, 1H), 8.02 (s, 1H), 7.68 (d, 1H),  
7.53 (d, 2H), 7.45 (m, 2H), 7.31 (m, 2H), 7.23 (m, 1H), 7.08 (m, 4H), 7.08 (m,  
2H), 7.06 (m, 2H), 6.75 (s, 1H), 4.65 (m, 1H), 3.53 (d, 1H), 3.41 (d, 1H), 3.04 (m,  
2H), 3.00 (m, 1H), 2.71 (m, 2H), 2.30 (m, 2H), 2.03 (m, 2H), 1.72 (m, 4H), 1.68  
25 (m, 2H), 1.27 (m, 1H)

Example 55:

Preparation of

30 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-[N-(2-fluorobenzoyl)amino]phenyl]piperidin-3-one

In accordance with the same procedure as in Example 48, except that

2-fluorobenzoyl chloride was used instead of ethyl chloroformate in Step 1 thereof, 220 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 9.91 (d, 1H), 8.63 (d, 1H), 8.18 (t, 1H), 7.79 (t, 1H), 7.43 (m, 1H), 7.21 (m, 4H), 7.20 (m, 4H), 7.04 (m, 2H), 6.68 (s, 1H), 5.27 (m, 1H), 4.13 (d, 1H), 3.31 (m, 1H), 3.00 (m, 2H), 2.87 (m, 1H), 2.34 (m, 2H), 2.02 (m, 1H), 2.00 (m, 4H), 1.67 (m, 1H), 1.64 (m, 3H), 1.22 (m, 1H)

#### Example 56:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-[N-(acetyl)amino]phenyl]piperidin-3-one

In accordance with the same procedure as in Example 48, except that acetyl chloride was used instead of ethyl chloroformate in Step 1 thereof, 190 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.31 (d, 1H), 7.70 (d, 1H), 7.56 (m, 1H), 7.54 (m, 2H), 7.33 (m, 1H), 7.15 (m, 2H), 7.23 (m, 1H), 7.08 (m, 4H), 7.08 (m, 2H), 7.06 (m, 3H), 6.65 (s, 1H), 4.99 (m, 1H), 4.37 (m, 1H), 3.16 (m, 1H), 2.81 (m, 1H), 2.79 (m, 2H), 2.27 (m, 3H), 2.25 (s, 3H), 2.00 (m, 2H), 1.74 (m, 4H), 1.67 (m, 2H), 1.31 (m, 1H)

#### Example 57:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-[N-(2,5-dihydropyrrol-1-yl)amino]phenyl]piperidin-3-one

Step 1.

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-[N-(2,5-dihydropyrrol-1-yl)amino]phenyl]piperidin-3-ol

To the solution of cis-2-buten-1,4-diol diacetate (1.15 g) in 30 ml of dichloromethane, was added the solution of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-aminophenyl)piperidin-3-ol (1.5 g) obtained in Step 2 of Example 43 in 20 ml of dimethylformamide. 0.44 ml of triethylamine was added to the reaction mixture, which was then stirred overnight at room temperature. 20 ml of ice water was added to the reaction mixture. The resulting organic layer was dried and concentrated on sodium sulfate. The resulting residue was purified with column chromatography to give 0.7 g of the titled compound.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.80-7.00 (m, 6H), 7.7 (m, 1H), 6.17 (t, 2H), 4.8 (m, 1H), 3.78 (s, 2H), 3.67 (s, 2H), 3.45 (m, 3H), 3.30 (m, 1H), 3.00 (m, 1H), 2.69 (m, 1H), 2.33 (d, 2H), 2.01 (t, 2H), 1.80-1.68 (m, 2H), 1.43 (t, 2H), 1.26 (d, 2H)

#### Step 2.

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-[N-(2,5-dihydropyrrol-1-yl)amino]phenyl]piperidin-3-one

To the solution of 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-[N-(2,5-dihydropyrrol-1-yl)amino]phenyl]piperidin-3-ol (0.7 g) obtained in Step 1 in 6 ml of dimethylsulfoxide (DMSO), was added 0.8 ml of triethylamine. The reaction mixture was cooled to 0°C. Pyridine-SO<sub>3</sub> complex (0.75 g) was added slowly to the reaction mixture, which was then stirred for about 2 hours at room temperature. The reaction mixture was diluted with 50 ml of ethyl acetate and washed with brine. The resulting organic layer was dried and concentrated on sodium sulfate. The resulting residue was filtered and dried to give 0.6 g of the titled compound as white color.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.70 (d, 1H), 7.54 (d, 1H), 7.49-7.40 (m, 2H), 7.32-7.22 (m, 4H), 7.09 (t, 1H), 7.01 (d, 1H), 4.58 (m, 1H), 3.73-3.70 (m, 1H), 3.57-3.54 (m, 1H), 3.49-3.45 (m, 4H), 2.91-2.85 (m, 2H), 2.50 (m, 1H), 2.32-2.28 (m, 2H), 1.98 (t, 2H), 1.77-1.55 (m, 7H), 1.35 (t, 1H)

#### Example 58:

## Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(tetrazol-1-yl)phenyl]piperidin-3-one

## 5 Step 1.

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(tetrazol-1-yl)phenyl]piperidin-3-ol

The mixture of  
10 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-aminophenyl)piperidin-3-one (2 g, 4 mmol) obtained in Step 2 of Example 43, sodium azide ( $\text{NaN}_3$ ; 409 mg, 6.2 mmol) and triethyl orthoformate (1.1 ml, 6.7 mmol) was refluxed in 10 ml of acetic acid for about 4 hours. The reaction mixture was cooled to room temperature. 10 ml of ice water was added to the  
15 reaction mixture, which was then stirred for about 48 hours. The resulting residue was filtered and dried to give 0.6 g of the titled compound.

$^1\text{H-NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  9.39 (s, 1H), 7.62 (d, 1H), 7.43 (m, 3H), 7.22 (m, 2H), 7.19 (m, 3H), 7.11 (m, 2H), 4.26(s, 1H), 3.87 (m, 1H), 3.65 (m, 1H), 3.21 (m, 1H), 2.66 (m, 1H), 2.45 (m, 2H), 2.21 (m, 2H), 2.05 (m, 3H), 1.74  
20 (m, 4H), 1.54 (m, 2H), 1.23 (m, 1H)

## Step 2.

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(tetrazol-1-yl)phenyl]piperidin-3-one

25

To the solution of 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(tetrazol-1-yl)phenyl]piperidin-3-ol (0.6 g) obtained in Step 1 in 10 ml of dimethylsulfoxide, was added 0.48 ml of triethylamine (3.4 mmol). The reaction mixture was cooled to 0°C.  
30 Pyridine- $\text{SO}_3$  complex (0.54 g, 3.4 mmol) was added to the reaction mixture, which was then stirred for about 2 hours at room temperature. The reaction mixture was diluted with 50 ml of ethyl acetate and washed with brine. The



resulting organic layer was dried and concentrated on sodium sulfate. The resulting residue was filtered and dried to give 0.4 g of the titled compound.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 9.08 (s, 1H), 7.67 (d, 1H), 7.53 (m, 2H), 7.49 (m, 1H), 7.33 (m, 2H), 7.25 (d, 2H), 6.76 (s, 1H), 4.56 (m, 1H), 3.60 (d, 1H), 3.48 (d, 1H), 3.05 (t, 1H) 2.94 (m, 1H), 2.35 (m, 1H), 2.29 (m, 2H), 1.97 (m, 2H), 1.75 (m, 4H), 1.52 (m, 3H) , 1.49 (m, 1H)

#### Example 59:

##### Preparation of

10 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(pyrrol-1-yl)-5-fluorophenyl]piperidin-3-one

##### Step 1.

15 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-amino-5-fluorophenyl)piperidin-3-ol

To the solution of 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-3-piperidinol hydrochloride (6.0 g) obtained in Step 2 of Example 1 in 50 ml of ethanol, were added 3.7 ml 2,5-difluoronitrobenzene and 4.8 ml triethylamine. The reaction mixture was refluxed overnight and concentrated. The resulting residue was filtered and dried to give 7.2 g of the titled compound.

##### Step 2.

25 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(pyrrol-1-yl)-5-fluorophenyl]piperidin-3-ol

The mixture of 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-amino-5-fluorophenyl)piperidin-3-ol (2 g) obtained in Step 1 and 0.96 ml of 2,5-dimethoxytetrahydrofuran was refluxed for 1 hour and concentrated under reduced pressure. The resulting residue was diluted with 70 ml of dichloromethane, washed with a saturated sodium

bicarbonate solution and brine. The resulting organic layer was dried and concentrated on sodium sulfate. The resulting residue was purified with column chromatography to give 1.8 g of the titled compound as white color.

5 Step 3.

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(pyrrol-1-yl)-5-fluorophenyl]piperidin-3-one

To the solution 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(pyrrol-1-yl)-5-fluorophenyl]piperidin-3-ol (1.8 g) obtained in Step 1 in 10 ml of dimethylsulfoxide, was added 2 ml of triethylamine. The reaction mixture was cooled to 0°C. Pyridine-SO<sub>3</sub> complex (1.8 g) was added to the reaction mixture, which was then stirred for about 2 hours at room temperature. The reaction mixture was diluted with 50 ml of ethyl acetate, washed with brine. The resulting organic layer was dried and concentrated on sodium sulfate. The resulting residue was purified with column chromatography to give 1.5 g of the titled compound.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.70 (d, 1H), 7.54 (d, 1H), 7.49 (q, 2H), 7.32 (d, 1H), 7.29 (m, 1H), 7.17 (m, 1H), 6.85 (d, 1H), 6.69 (t, 1H), 6.27 (s, 1), 4.58 (t, 1H), 3.44 (q, 2H), 2.87 (m, 2H), 2.31 (m, 1H), 2.28 (d, 2H), 1.98 (m, 3H), 1.60-1.22 (m, 7H)

Example 60:

Preparation of

25 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(pyrrol-1-yl)phenyl]piperidin-3-one

In accordance with the same procedure as in Example 59, except that 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-aminophenyl)piperidin-3-ol obtained in Step 2 of Example 43 was used instead of 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-a

mino-5-fluorophenyl)piperidin-3-ol in Step 1 of Example 59, 1.3 g of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.86 (d, 1H), 7.67 (d, 1H), 7.53 (m, 2H), 7.45 (m, 2H), 7.24 (m, 2H), 7.01 (m, 1H), 6.99 (m, 1H), 6.27 (s, 1H), 4.59 (m, 1H), 3.48 (m, 2H), 2.99 (m, 2H), 2.50 (m, 1H), 2.25 (m, 2H), 1.91 (m, 4H), 1.42 (m, 4H)

#### Example 61:

##### Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(N,N-dimethylcarbamoyl)phenyl]piperidin-3-one

##### Step 1.

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-formylphenyl)piperidin-3-ol

In accordance with the same procedure as in Step 3 of Example 1, except that 2-fluorobenzaldehyde was used instead of 1-fluorophenyl methyl sulfone, 5.3 g of the titled compound was prepared.

Step 2. 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(N,N-dimethylcarbamoyl)phenyl]piperidin-3-ol

To the solution of 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-formylphenyl)piperidin-3-ol (4.0 g) obtained in Step 1 and sodium hydrogen phosphate (NaH<sub>2</sub>PO<sub>4</sub>, 340 mg) in the mixed solvent of 20 ml of acetonitrile and 6 ml of water, were added 1.0 ml of 35%-hydrogen peroxide solution and 30 ml of sodium hypochlorite solution (NaClO<sub>4</sub>) at 10 °C. The reaction mixture was stirred for about 1 hour at room temperature. Sodium sulfite (1.0 g) was added to the reaction mixture, which was then acidified with 10%-hydrochloride solution. The resulting solid was filtered and dried to give 3.6 g of a solid.

The solution of the resulting solid (2.5 g), dimethylamine hydrochloride (1.0 g) and benzotriazol-1-yloxytripyrrolidinophosphonium hexaphosphate (PyBOP, 3.1 g) in 30 ml of dichloromethane was cooled to 0°C. Diisopropylethylamine (DIEA, 1.0 g) was added to the reaction mixture, which was then stirred for about 4 hours at room temperature. 30 ml of water was added to the reaction mixture. The resulting organic layer was dried and concentrated on sodium sulfate. The resulting residue was purified with column chromatography to give 1.5 g of the titled compound.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.69 (d, 1H), 7.57 (m, 2H), 7.49 (m, 2H), 7.31 (m, 1H), 7.06 (t, 1H), 6.97 (d, 1H), 6.73 (s, 1H), 3.78 (m, 1H), 3.64 (m, 2H), 3.41 (m, 2H), 3.01 (s, 3H), 2.78 (m, 4H), 2.31 (broad, 2H), 2.00 (m, 3H), 1.75 (m, 2H), 1.52 (m, 2H), 1.38 (m, 1H)

### Step 3.

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(N,N-dimethylcarbamoyl)phenyl]piperidin-3-one

To the solution of 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(N,N-dimethylcarbamoyl)phenyl]piperidin-3-ol (1.8 g) obtained in Step 2 in 10 ml of dimethylsulfoxide (DMSO), was added 2 ml of triethylamine. The reaction mixture was cooled to 0°C. Pyridine-SO<sub>3</sub> complex (1.8 g) was added slowly to the reaction mixture, which was then stirred for about 2 hours at room temperature. The reaction mixture was diluted with 50 ml of ethyl acetate, washed with brine. The resulting organic layer was dried and concentrated on sodium sulfate. The resulting residue was purified with column chromatography to give 1.2 g of the titled compound.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.69 (d, 1H), 7.56-7.45 (m, 4H), 7.34-7.23 (m, 3H), 7.07 (s, 1H), 6.97 (t, 1H), 6.64 (s, 1H), 4.63 (s, 1H), 3.79-3.29 (m, 4H), 3.07 (s, 1H), 2.77 (s, 3H), 2.64 (s, 1H), 2.30 (d, 2H), 2.01 (t, 2H), 1.75-1.69 (m, 5H), 1.43 (d, 2H), 1.22 (m, 2H)

Example 62:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(N  
5 -methylcarbamoyl)phenyl]piperidin-3-one

In accordance with the same procedure as in Example 61, except methylamine was used instead of dimethylamine hydrochloride in Step 1 thereof, 1.05 g of the titled compound was prepared.

10  $^1\text{H-NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  8.56 (broad, 1H), 8.05 (d, 1H), 7.70 (d, 1H), 7.54 (d, 1H), 7.49-7.22 (m, 7H), 7.12 (t, 1H), 6.75 (s, 1H), 4.62 (t, 1H), 3.72 (m, 1H), 3.66 (s, 2H), 3.27 (q, 2H), 2.96 (s, 3H), 2.78 (m, 2H), 2.31 (d, 2H), 1.99 (t, 2H), 1.78 (m, 5H), 1.55 (m, 2H), 1.22 (m, 2H)

15 Example 63:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-carboxyphenyl)piperidin-3-one

20 To the solution of 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-formylphenyl)piperidin-3-one (300 mg, 0.62 mmol) obtained in Example 15 and sodium hydrogen phosphate (22 mg,  $\text{NaH}_2\text{PO}_4$ ) in the mixed solvent of 1 ml of acetonitrile and 0.5 ml of water, was added 35%-hydrogen peroxide solution (0.1 ml) and 3 ml of sodium  
25 hypochlorite solution ( $\text{NaClO}_4$ ; 78mg) at 10°C. The reaction mixture was stirred for about 1 hour at room temperature. Sodium sulfite (1.0 g) was added to the reaction mixture, which was then acidified with 10%-hydrochloride solution. The resulting solid was filtered and dried to give 150 mg of the titled compound.

30  $^1\text{H-NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  7.55-6.74 (m, 9H), 4.63 (m, 1H), 3.73-3.59 (m, 4H), 3.51-3.45 (m, 2H), 2.78 (m, 2H), 2.31 (d, 2H), 1.99 (t, 2H), 1.78 (m, 5H), 1.55 (m, 2H), 1.22 (m, 2H)

Example 64:

## Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-carboxyphenyl)piperidin-3-one

In accordance with the same procedure as in Example 63, except 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-formylphenyl)piperidin-3-one obtained in Example 14 was used instead of 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-formylphenyl)piperidin-3-one, 130 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.04 (d, 1H), 7.67 (d, 1H), 7.50 (m, 3H), 7.28 (m, 3H), 6.95 (s, 1H), 4.57 (m, 1H), 4.55-3.90 (m, 4H), 2.27 (m, 2H), 2.20 (m, 2H), 1.98 (m, 2H), 1.55 (m, 2H), 1.20 (m, 1H)

Example 65:

## Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(1,3-oxazol-5-yl)phenyl]piperidin-3-one

## Step 1.

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(1,3-oxazol-5-yl)phenyl]piperidin-3-ol

The mixture of 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-formylphenyl)piperidin-3-ol (3 g, 6 mmol) obtained in Step 1 of Example 61, potassium carbonate (2 g, 20 mmol) and p-toluenesulfonylmethyl isocyanate (TosMIC; 1.8 g, 9 mmol) was refluxed in 500 ml of methanol for about 2 hours. 150 ml of water was added to the concentrated reaction mixture, which was then extracted with dichloromethane.

The resulting organic layer was dried and concentrated on sodium sulfate. The resulting residue was purified with column chromatography to give 3.2 g of

the titled compound.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.89 (s, 1H), 7.71 (m, 3H), 7.54 (m, 1H), 7.32 (m, 1H), 7.26 (m, 1H), 7.22 (m, 2H), 6.88 (m, 1H), 6.82 (s, 1H), 4.16 (s, 1H), 3.85 (m, 1H), 3.69 (m, 1H), 3.34 (m, 1H), 3.06 (m, 1H), 2.73 (m, 2H), 2.28 (m, 2H), 2.00 (m, 3H), 1.79 (m, 4H), 1.54 (m, 2H), 1.32 (m, 1H)

Step 2.

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(1, 3-oxazol-5-yl)phenyl]piperidin-3-one

To the solution of 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(1, 3-oxazol-5-yl)phenyl]piperidin-3-ol (3.36 g, 6 mmol) obtained in Step 1 in 20 ml of dimethyl sulfoxide (DMSO), was added 2.7 ml of triethylamine (19 mmol). The reaction mixture was cooled to 0°C. Pyridine-SO<sub>3</sub> complex (3.04 g, 19 mmol) was added to the reaction mixture, which was then stirred for about 2 hours at room temperature. The reaction mixture was diluted with 100 ml of ethyl acetate, washed with brine. The resulting organic layer was dried and concentrated on sodium sulfate. The resulting residue was purified with column chromatography to give 1.67 g of the titled compound.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.75 (s, 1H), 7.89 (d, 1H), 7.75 (d, 1H), 7.60 (m, 3H), 7.41 (m, 2H), 7.33 (m, 2H), 7.19 (m, 2H), 6.73 (s, 1H), 4.65 (m, 1H), 3.65 (q, 2H), 3.26 (m, 2H), 3.70 (m, 1H), 2.36 (m, 2H), 2.02 (m, 2H), 1.79 (m, 4H), 1.55 (m, 2H)

#### Example 66:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-hydroxyphenyl)piperidin-3-one

Step 1.

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-hy

droxyphenyl)piperidin-3-ol

To the solution of  
4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-hydroxyphenyl)piperidin-3-ol (1.38 g, 2.8 mmol) obtained in Step 1 of Example 61 in  
5 50 ml of chloroform, was added 486 mg of m-chloroperbenzoic acid (5.6 mmol).

The reaction mixture was stirred for about 2 hours at room temperature. The resulting organic layer was washed with the solutions of sulfurous acid and sodium bicarbonate, dried on sodium sulfate. The resulting ester residue was  
10 dissolved in 100 ml of methanol and was added a drop of concentrated hydrochloride thereto. The reaction mixture was stirred for about 1 hour, was added sodium bicarbonate (0.5 g) thereto. The reaction mixture was concentrated. The resulting residue was purified with column chromatography to give 1.0 g of the titled compound.

15 Step 2.

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-hydroxyphenyl)phenyl]piperidin-3-one

20 The solution of  
4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-hydroxyphenyl)piperidin-3-ol (200 mg, 0.41 mmol) obtained in Step 1 in 10 ml of dimethyl sulfoxide (DMSO) was cooled to 0°C and was added 199 mg of pyridine-SO<sub>3</sub> complex (1.25 mmol) thereto. The reaction mixture was stirred  
25 for about 2 hours at room temperature, diluted with 100 ml of ethyl acetate, and washed with brine. The resulting organic layer was dried and concentrated on sodium sulfate. The resulting residue was purified with column chromatography to give 120 mg of the titled compound.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.68 (d, 1H), 7.54 (d, 1H), 7.53 (m, 6H),  
30 7.42 (m, 1H), 7.01 (m, 1H), 6.86 (d, 1H), 4.81 (m, 1H), 3.91 (t, 1H), 3.45 (m, 2H), 2.92 (m, 1H), 2.35 (m, 3H) 2.2.04 (m, 3H), 2.35 (m, 2H), 2.07 (m, 2H), 1.65 (m, 4H), 1.52 (m, 2H), 1.35 (m, 1H)



Example 67:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(1  
5 H-tetrazol-5-yl)phenyl]piperidin-3-one

Step1.

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-cy  
anophenyl)piperidine-3-ol

10

In accordance with the same procedure as in Step 3 of Example 1, except that 2-fluorobenzonitrile was used instead of 1-fluorophenyl methyl sulfone, 2.5 g of the titled compound was prepared.

15 Step 2.

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(1  
H-tetrazol-5-yl)phenyl]piperidin-3-ol

To the solution of  
20 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-cy  
anophenyl)piperidin-3-ol (700 mg, 1.43 mmol) obtained in Step 1 in 20 ml of  
dimethylformamide, were added ammonium chloride (769 mg) and sodium  
azide (934 mg). The reaction mixture was stirred for about 12 hours at 120 °C.

20 ml of water was added to the reaction mixture, which was then acidified  
25 with a concentrated hydrochloride solution. The resulting solid was filtered  
and dissolved in dichloromethane. The resulting residue was purified with  
column chromatography to give 120 mg of the titled compound.

Step 3.

30 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(1  
H-tetrazol-5-yl)phenyl]piperidin-3-one

To the solution of 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(1 H-tetrazol-5-yl)phenyl]piperidin-3-ol (120 mg, 0.22 mmol) obtained in Step 2 in 5 ml of dimethyl sulfoxide (DMSO), was added 0.093 ml of triethylamine (0.66 mmol). The reaction mixture was cooled to 0°C. Pyridine-SO<sub>3</sub> complex (103 mg, 0.66 mmol) was added to the reaction mixture, which was then stirred for about 2 hours at room temperature. The reaction mixture was diluted with 100 ml of ethyl acetate, washed with brine. The resulting organic layer was dried on sodium sulfate. The resulting residue was purified with column chromatography to give 50 mg of the titled compound.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.75-7.64 (m, 4H), 7.53-7.44 (m, 9H), 7.42 (d, 1H), 7.35-7.26 (m, 2H), 6.74 (s, 1H), 4.56 (m, 1H), 4.00 (q, 1H), 3.86 (q, 1H), 2.89 (m, 1H), 2.34 (m, 2H), 2.24-2.18 (m, 3H), 2.02 (m, 2H), 1.91 (m, 1H), 1.75 (m, 2H), 1.54 (t, 2H), 1.40 (m, 1H)

#### Example 68:

##### Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methylsulfinylphenyl)phenyl]piperidin-3-one

The solution of 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methylthiophenyl)piperidin-3-one (68 mg, 0.13 mmol) obtained in Example 36 in 2 ml of dichloromethane was cooled to -78°C. 30 mg of 70% 3-chloroperbenzoic acid was added to the reaction mixture, which was then stirred for about 1 hour at room temperature. The resulting organic layer was washed with a saturated sodium bicarbonate solution, dried and concentrated on sodium sulfate. The resulting residue was purified with column chromatography to give 30 mg of the titled compound.

R<sub>f</sub> = 0.1 (CH<sub>2</sub>Cl<sub>2</sub>/ether=1:1)

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.92 (d, 1H), 7.69 (d, 1H), 7.42 (m, 6H), 7.29 (m, 1H), 7.15 (m, 1H), 6.78 (d, 1H), 4.65 (m, 1H), 3.75 (t, 1H), 3.45 (m,

2H), 3.22 (m, 1H), 2.75 (s, 1H) 2.71 (s; 3H), 2.35 (m, 2H), 2.07 (m, 2H), 1.65 (m, 4H), 1.52 (m, 2H), 1.35 (m, 1H)

Example 69:

5 Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-ethylsulfinylphenyl)phenyl]piperidin-3-one

10 The solution of 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-ethylthiophenyl)piperidin-3-one (38 mg) obtained in Example 42 was cooled to -78°C. 70% 3-chloroperbenzoic acid (20 mg) was added to the reaction mixture, which was then stirred for about 1 hour at room temperature. The resulting organic layer was washed with a saturated sodium bicarbonate solution, dried and concentrated on sodium sulfate. The resulting residue was purified with column chromatography to give 22 mg of the titled compound.

15 <sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.87 (d, 1H), 7.74 (d, 1H), 7.53 (m, 6H), 7.33 (m, 3H), 7.15 (dd, 1H), 6.73 (d, 1H), 4.65 (m, 1H), 3.69 (d, 1H), 3.55 (d, 1H), 3.45 (d, 1H), 3.15 (t, 1H), 2.89 (q, 2H), 2.71 (m, 1H), 2.31 (d, 2H), 2.05 (t, 20 2H), 1.89 (m, 1H), 1.75 (m, 3H), 1.54 (m, 2H), 1.36 (m, 1H) 1.34 (t, 3H)

Example 70:

Preparation of

25 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-isopropylsulfinylphenyl)piperidin-3-one

Step 1.

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-isopropylthiophenyl)piperidin-3-ol

30

In accordance with the same procedure as in Step 1 of Example 24, except that 1-bromo-2-isopropylthiobenzene was used instead of

1-bromo-4-fluoro-2-trifluoromethylbenzene, 66 mg of the titled compound was prepared.

Step 2.

- 5 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-isopropylsulfinylphenyl)piperidin-3-one

To the solution of 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-isopropylthiophenyl)piperidin-3-ol (60 mg) obtained in Step 1 in 5 ml of dimethyl sulfoxide (DMSO), was added 0.093 ml of  
10 triethylamine (0.66 mmol). The reaction mixture was cooled to 0°C. Pyridine-SO<sub>3</sub> complex (103 mg, 0.66 mmol) was added to the reaction mixture, which was then stirred for about 2 hours at room temperature. The reaction mixture was diluted with 100 ml of ethyl acetate, washed with brine. The resulting organic layer was dried on sodium sulfate. The resulting residue was  
15 purified with column chromatography to give 50 mg of the titled compound.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.70 (d, 1H), 7.60-7.40 (m, 4H), 7.33 (m, 1H), 7.25 (m, 2H), 7.15 (m, 1H), 7.05 (m, 1H), 6.98 (d, 1H), 6.73 (s, 1H), 4.66 (m, 1H), 3.71 (d, 1H), 3.61 (d, 1H), 3.51 (m, 1H), 3.43 (m, 1H), 3.15 (t, 1H), 2.71 (m, 1H), 2.33 (d, 2H), 1.98 (t, 2H), 1.85 (m, 1H), 1.72 (m, 3H), 1.55 (d, 3H),  
20 1.53 (m, 2H), 1.35 (m, 1H), 1.26 (d, 3H)

Example 71:

Preparation of

- 25 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-N-(acryloyl)aminophenyl]piperidin-3-one

In accordance with the same procedure as in Example 48, except acryloyl chloride was used instead of ethyl chloroformate in Step 1 thereof, 254 mg of the titled compound was prepared.

30 <sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.30 (s, 1H), 7.62 (d, 1H), 7.40-7.50(m, 3H), 6.41 (d, 1H), 5.83 (d, 1H), 4.70 (m, 1H), 3.64 (dd, 2H), 2.00-1.30 (m, 10H)

Example 72:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(3,3-dimethylureido)phenyl]piperidin-3-one

5

In accordance with the same procedure as in Example 48, except dimethylcarbamoyl chloride was used instead of ethyl chloroformate in Step 1 thereof, 220 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.10 (d,1H), 7.60-7.50 (m, 4H), 7.52-7.10 (m, 5H), 6.72 (s,1H), 4.70 (m, 1H), 3.80-3.12 (m, 3H), 2.22-1.90 (m, 2H), 1.90-1.32 (m, 11H)

10

Example 73:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(methoxycarbonyl)phenyl]piperidin-3-one

15

In accordance with the same procedure as in Example 24, except methyl 2-bromobenzoate was used instead of

1-bromo-4-fluoro-2-trifluoromethyl benzene in Step 1 thereof, 70 mg of the titled compound was prepared.

20

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.80 (m, 2H), 7.62 (m,1H), 7.40-7.20 (m, 4H), 6.72 (s, 1H), 4.73 (m, 1H), 3.80 (s, 3H), 2.63 (m, 1H), 2.00-1.32 (m, 10H)

Example 74:

25

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-phenylpiperidin-3-one

In accordance with the same procedure as in Example 24, except methyl bromobenzene was used instead of 1-bromo-4-fluoro-2-trifluoromethyl benzene in Step 1 thereof, 35 mg of the titled compound was prepared.

30

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.57 (m, 6H), 7.31 (m, 2H), 6.89 (m, 1H), 6.45 (s, 1H), 7.05 (d, 1H), 6.73 (s, 1H), 4.66 (m, 1H), 4.22 (m, 1H), 4.05 (d, 1H), 3.75 (d, 1H), 3.53 (m, 1H), 3.44 (m, 1H), 2.75 (m, 1H), 2.31 (d, 1H), 2.05 (t, 2H), 1.89 (m, 1H), 1.75 (m, 3H), 1.54 (m, 2H), 1.36 (m, 1H)

5

Example 75:

Preparation of

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(1,1-dioxo-isothiazolin-2-yl)phenyl]]piperidin-3-one

10

In accordance with the same procedure as in Example 43, except 3-chloropropanesulfonyl chloride was used instead of 3-chloropivaloyl chloride in Step 3 thereof, 65 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) : 7.78 (d, 1H), 7.69 (m, 2H), 7.56 (m, 2H), 7.46 (m, 1H), 7.25 (m, 1H), 7.05 (m, 2H), 6.71 (s, 1H), 4.65 (m, 1H), 3.94 (d, 1H), 3.85 (m, 1H), 3.58 (m, 1H), 3.51 (m, 1H), 3.26 (m, 3H), 2.76 (m, 1H), 2.48 (m, 2H), 2.33 (m, 2H), 2.04 (m, 3H), 1.78 (m, 4H), 1.53 (m, 3H), 1.26 (m, 1H)

15

Example 76:

20 Preparation of

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

Step 1. Methyl 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylate

25

The solution of methyl 1-amino-1-cyclohexanecarboxylate (11.3 g, 58.5 mmol) and 2-furan-carboxylic acid (5.0g) in 100 ml of dimethylformamide was cooled to 0 °C. To the reaction mixture, were added diisopropylethylamine (29 ml, 175.5 mmol), 1-hydroxybenzotriazole hydrate (9.1g, 67.5 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC-HCl; 12.9 g, 67.5 mmol). The reaction mixture was stirred for 24 hours at room temperature. 50 g of ice water was added to the reaction mixture to stop the

30

reaction. The reaction mixture was diluted with 200 ml of ethyl acetate; washed with 10% citric acid solution and a saturated sodium bicarbonate solution, and then dried and concentrated on sodium sulfate to give 9.6 g of the titled compound. The titled compound was used in the next-step without  
5 further purifications

Step 2. 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid

To the solution of methyl  
10 1-[N-(furan-2-ylcarbonyl)amino]-1-cyclohexanecarboxylate (9.6 g) obtained in Step 1 in 100 ml of methanol, was added 2N-sodium hydroxide solution (50 ml).

The reaction mixture was refluxed for 3 hours and concentrated. The resulting residue was diluted with 150 ml of water and washed with 100 ml of diethyl ether. 6N-hydrochloride was added dropwise to the resulting aqueous  
15 layer to adjust the pH of the solution to pH 3. The resulting white solid was filtered and dried to give the titled compound (6.1 g).

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 2.15 (d, 2H), 1.97 (t, 2H), 1.70 (m, 3H), 1.49-1.37 (m, 3H)

20 Step 3.

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-tert-butoxy carbonyl-3-piperidinol

To the solution of tert-butyl 4-amino-3-hydroxypiperidin-1-carboxylate (20  
25 g, 92 mmol) obtained in step 4 of Preparation Example 1 and 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid (5.0 g) obtained in Step 2 in 70 ml of dimethylformamide, were added hydroxybenzotriazole(HOBt; 3.1 g, 23.1 mmol) and diisopropylethylamine (DIEA; 9.0 ml, 53.4 mmol). The reaction mixture was cooled to 0°C and was added  
30 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC-HCl; 5.1 g, 26.7 mmol) thereto. The reaction mixture was stirred for 4 hours at room temperature. The reaction mixture was diluted with 200 ml of ethyl acetate,

washed with 10% citric acid solution, a saturated sodium bicarbonate solution and brine. The resulting organic layer was dried and concentrated on sodium sulfate. The resulting residue was purified with column chromatography to give 4.3 g of the titled compound.

5

Step 4.

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-3-piperidinol hydrochloride

10 The solution of 4-{1-[N-(furan-2-ylcarbonyl)amino]cyclohexane carbonyl}amino-1-tert-butoxycarbonyl-3-piperidinol (30g) obtained in Step 3 in 100ml of 3N-hydrochloride/ethyl acetate was stirred for about 2 hours at room temperature. 100ml of diethyl ether was added to the reaction mixture, and then the resulting white solid was filtered and dried to give 3.6 g of the titled  
15 compound.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.84 (d, 1H), 7.61 (s, 1H), 7.31 (d, 1H), 7.16 (d, 1H), 6.63 (s, 1H), 3.38 (m, 1H), 3.14 (m, 1H), 2.88 (d, 1H), 2.73 (d, 1H), 2.33 (t, 1H), 2.15 (m, 2H), 1.73 (m, 3H), 1.41 (m, 2H), 1.31 (m, 1H)

20 Step 5.

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methane sulfonylphenyl)piperidin-3-ol

To the solution of 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]  
25 cyclohexanecarbonyl]amino]-3-piperidinol hydrochloride (600 mg, 1.55 mmol) obtained in Step 4 and 1-flourophenyl methyl sulfone (272 g, 1.55 mmol) in 30 ml of dimethylformamide, was added potassium carbonate (500 mg). The reaction mixture was stirred overnight at about 100°C and cooled to room temperature. The reaction mixture was diluted with 200ml of ethyl acetate and  
30 washed with brine. The resulting organic layer was dried and concentrated on sodium sulfate. The resulting residue was purified with column chromatography to give 390 mg of the titled compound as white solid.



<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.78 (d, 1H), 7.50 (s, 1H), 7.43 (m, 1H), 7.31 (m, 1H), 7.13 (m, 2H), 6.55 (m, 2H), 3.88 (m, 1H), 3.69 (m, 1H), 3.27 (m, 5H), 2.81 (m, 1H), 2.67 (t, 1H), 2.22 (t, 2H), 1.99 (m, 3H), 1.70 (m, 4H), 1.48-1.29 (m, 3H)

5

Step 6.

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

10 To the solution of 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-ol (0.39g, 0.77 mmol) obtained in Step 5 in 3 ml of dimethylsulfoxide (DMSO), was added 0.4 ml of triethylamine. The reaction mixture was cooled to 0°C. Pyridine-SO<sub>3</sub> complex (0.4 g, 2.51 mmol) was  
15 added to the reaction mixture, which was then stirred for 2 hours. The reaction mixture was diluted with 20 ml of ethyl acetate, washed with brine. The resulting organic layer was dried and concentrated on sodium sulfate. The resulting residue was purified with column chromatography to give 155 mg of the titled compound as white solid.

20 <sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.1 (d, 1H), 7.6 (m, 2H), 6.5 (m, 1H), 6.4 (s, 1H), 4.7 (m, 1H), 3.7 (dd, 2H), 3.2 (s, 3H), 2.4 (m, 2H), 1.4-2.0 (m, 10H)

#### Example 77:

Preparation of

25 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(3-fluoro-2-methylsulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that 2,6-difluorophenyl methyl sulfone was used instead of 1-fluorophenyl methyl sulfone in Step 5 thereof, 230 mg of the titled compound was prepared.  
30

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.48 (m, 1H), 7.46 (m, 2H), 7.13 (d, 1H),

7.06 (m, 2H), 6.53 (d, 1H), 6.44 (s, 1H), 4.68 (m, 1H), 3.73 (m, 2H), 3.68 (d, 1H), 3.25 (s, 3H), 2.66 (m, 1H), 2.24 (d, 2H), 2.00 (t, 3H), 1.75 (m, 3H), 1.63 (m, 2H), 1.47 (m, 1H)

5 Example 78:

Preparation of

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-nitrophenyl)piperidin-3-one

10 In accordance with the same procedure as in Example 76, except that 2,5-difluoronitrobenzene was used instead of 1-fluorophenyl methyl sulfone in Step 5 thereof, 170 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.50-7.22 (m, 5H), 6.55 (d, 1H), 6.32 (s, 1H), 4.70 (m, 1H), 3.86 (q, 3H), 3.41 (t, 1H), 2.78 (m, 1H), 2.31 (broad, 2H),  
15 2.00 (m, 3H), 1.75 (m, 2H), 1.52 (m, 2H), 1.38 (m, 1H)

Example 79:

Preparation of

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-formylphenyl)piperidin-3-one  
20

In accordance with the same procedure as in Example 76, except that 2,5-difluorobenzaldehyde was used instead of 1-fluorophenyl methyl sulfone in Step 5 thereof, 190 mg of the titled compound was prepared.

25 <sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 10.32 (s, 1H), 7.77 (d, 1H), 7.52-7.42 (m, 2H), 7.15 (m, 2H), 6.53 (s, 1H), 4.65 (m, 1H), 3.58 (m, 2H), 3.34 (m, 1H), 3.14 (t, 1H), 2.70 (m, 1H), 2.38 (s, 3H), 2.23(m, 2H), 1.94 (m, 2H), 1.73-1.51 (m, 5H), 1.43-1.21 (m, 3H)

30 Example 80:

Preparation of

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-isopropy

## Isulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that 1-fluorophenyl isopropyl sulfone was used instead of 1-fluorophenyl methyl sulfone in Step 5 thereof, 173 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.06 (d, 1H), 7.49 (t, 1H), 7.38 (m, 2H), 7.26 (m, 2H), 7.15 (d, 1H), 6.53 (m, 1H), 6.44 (s, 1H), 6.65 (m, 1H), 3.80 (m, 2H), 3.74 (d, 2H), 3.15 (t, 1H), 2.67 (m, 1H), 2.17 (m, 2H), 1.98 (m, 3H), 1.72 (d, 3H), 1.50 (m, 2H), 1.47 (m, 2H), 1.31 (m, 1H), 1.29 (d, 3H), 1.09 (d, 3H)

Example 81:

Preparation of

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-phenylsulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that 2-fluorophenyl sulfone was used instead of 1-fluorophenyl methyl sulfone in Step 5 thereof, 150 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) : 7.63 (m, 1H), 7.50-7.15 (m, 10H), 6.71 (s, 1H), 4.38 (m, 1H), 3.67 (d, 1H), 3.46 (d, 1H), 3.34 (m, 1H), 3.21 (t, 1H), 2.63 (m, 1H), 2.26 (m, 2H), 1.94 (t, 2H), 1.71 (m, 3H), 1.47 (m, 3H), 1.26 (m, 3H)

Example 82:

Preparation of

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-methoxybenzylsulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that 2-fluorophenyl-(4-methoxy)benzyl sulfone was used instead of 1-fluorophenyl methyl sulfone in Step 5 thereof, 76 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) : 7.63 (m, 1H), 7.50-7.15 (m, 10H), 6.71 (s, 1H), 4.75 (m, 1H), 4.38 (m, 1H), 3.71 (s, 3H), 3.67 (d, 1H), 3.46 (d, 1H), 3.34 (m,

1H), 3.21 (t, 1H), 2.63 (m, 4H), 2.26 (m, 2H), 1.94 (t, 2H), 1.71 (m, 3H), 1.47 (m, 3H), 1.26 (m, 3H)

Example 83:

5 Preparation of

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-methylsulfonylphenyl)piperidin-3-one

10 In accordance with the same procedure as in Example 76, except that 2,5-difluorophenyl methyl sulfone was used instead of 1-fluorophenyl methyl sulfone in Step 5 thereof, 150 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.8 (m, 1H), 7.52 (m, 2H), 7.41 (m, 2H), 7.15 (s, 1H), 6.52 (s, 1H), 6.40 (s, 1H), 4.62 (s, 1H), 3.21 (s, 3H), 2.18-1.31 (m, 12H)

15

Example 84:

Preparation of

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-methoxy-2-methylsulfonylphenyl)piperidin-3-one

20

In accordance with the same procedure as in Example 76, except that 2-fluoro-5-methoxyphenyl methyl sulfone was used instead of 1-fluorophenyl methyl sulfone in Step 5 thereof, 135 mg of the titled compound was prepared.

25 <sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.83 (m, 1H), 7.52 (m, 2H), 7.41 (m, 2H), 7.18 (s, 1H), 6.54 (s, 1H), 6.38 (s, 1H), 4.65 (s, 1H), 3.85 (s, 3H), 3.21 (s, 3H), 2.2-1.15 (m, 12H)

Example 85:

Preparation of

30 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-thiomethoxyphenyl)piperidin-3-one

## Step 1.

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-thiomethoxyphenyl)piperidin-3-ol

5           To                           the                           suspension                           of  
4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-3-piperidinol  
hydrochloride (300 mg) obtained in Step 4 of Example 76 in 5 ml of toluene,  
were added 1-bromo-4-fluoro-2-thiomethoxybenzene (250 mg), cesium  
carbonate (1.72 g), (S)-2,2-bis(diphenylphosphino)-1,1-binaphthyl (BINAP; 150  
10 mg) and tris(dibenzylideneacetone)dipalladium (122 mg). The reaction  
mixture was stirred for about 8 hours at 100°C under nitrogen atmosphere.  
The reaction mixture was diluted with 30ml of ethyl acetate, filtered and  
concentrated. The resulting residue was purified with column chromatography  
to give 179 mg of the titled compound.

15           <sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.52 (m, 2H), 7.23 (m, 3H), 6.87 (d, 1H),  
6.53 (d, 2H), 3.81 (m, 1H), 3.62 (m, 1H), 3.38 (m, 1H), 3.19 (m, 1H), 2.70 (m,  
1H), 2.35 (s, 3H), 2.23(m, 2H), 1.94 (m, 2H), 1.73-1.51 (m, 5H), 1.43-1.21 (m,  
3H)

## 20   Step 2.

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-thiomethoxyphenyl)piperidin-3-one

                  To                           the                           solution                           of  
25   4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-thiomethoxyphenyl)piperidin-3-ol (0.15 g) obtained in Step 1 in 3 ml of  
dimethyl sulfoxide (DMSO), was added 0.16 ml of triethylamine. The reaction  
mixture was cooled to 0°C. Pyridine-SO<sub>3</sub> complex (0.15 g) was added to the  
reaction mixture, which was then stirred for about 4 hours at room temperature.  
30   The reaction mixture was diluted with 20 ml of ethyl acetate and washed with  
brine. The resulting organic layer was dried and concentrated. The resulting  
residue was purified with column chromatography to give 84 mg of the titled

compound as white solid.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.52 (s, 1H), 7.48 (d, 1H), 7.15 (d, 1H),  
6.99 (m, 1H), 6.77 (m, 2H), 6.52 (m, 1H), 6.45 (s, 1H), 4.67 (m, 1H), 3.64 (d,  
1H), 3.56 (d, 1H), 3.45 (m, 1H), 3.13 (m, 1H), 2.67 (m, 1H), 2.35 (s, 3H), 2.23  
5 (m, 2H), 2.00 (m, 2H), 1.97 (m, 1H), 1.73 (m, 6H), 1.51 (m, 2H), 1.44 (m, 1H)

Example 86:

Preparation of

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-cyanoph  
10 enyl)piperidin-3-one

In accordance with the same procedure as in Example 83, except that  
2-bromobenzonitrile was used instead of 1-bromo-4-fluoro-2-thiomethoxy  
benzene in Step 1 thereof, 160 mg of the titled compound was prepared.

15 <sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.50 (d, 1H), 7.48 (m, 3H), 7.25 (m, 1H),  
7.23 (d, 1H), 7.14 (t, 1H), 6.98 (d, 1H), 6.53 (q, 1H), 6.44 (s, 1H), 4.68 (m, 1H),  
3.89 (dt, 3H), 3.41 (t, 1H), 2.75 (m, 1H), 2.27 (m, 2H), 2.00-1.94 (m, 3H),  
1.73-1.69 (m, 3H), 1.43-1.21 (m, 3H)

20 Example 87:

Preparation of

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-  
methoxyacetaminophenyl)piperidin-3-one

25 Step 1.

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-  
nitrophenyl)piperidin-3-ol

To the solution of 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexane  
30 carbonyl]amino]-3-piperidinol hydrochloride (6.0 g) obtained in Step 4 of  
Example 76 in 50 ml of ethanol, were added 2,5-difluoronitrobenzene (3.7 ml)  
and triethylamine (4.8 ml). The reaction mixture was refluxed overnight and

concentrated. The resulting yellow solid was filtered and dried to give 7.2 g of the titled compound.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.52 (s, 1H), 7.48 (d, 1H), 7.15 (d, 1H), 6.99 (m, 1H), 6.77 (m, 2H), 6.52 (m, 1H), 6.45 (s, 1H), 4.67 (m, 1H), 3.64 (d, 1H), 3.56 (d, 1H), 3.45 (m, 1H), 3.13 (m, 1H), 2.77 (m, 2H), 2.23 (m, 2H), 2.00 (m, 2H), 1.97 (m, 1H), 1.73 (m, 6H), 1.51 (m, 2H), 1.44 (m, 1H)

#### Step 2.

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-aminophenyl)piperidine-3-ol

To the solution of 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-nitrophenyl)piperidin-3-ol (7.0 g, 14.7 mmol) obtained in Step 1 in 70 ml of ethanol, was added tin (II) chloride hydrate (1.13 mg, 44.2 mmol). The reaction mixture was refluxed for about 4 hours at 80 °C, and then cooled to room temperature and concentrated. The resulting residue was diluted with 100 ml of ethyl acetate, alkalized with 29% aqueous ammonia solution (pH 8), and then the resulting white solid was filtered out. 100 ml of distilled water was added to the resulting filtrate. The resulting organic layer was dried and concentrated on sodium sulfate to give the titled compound (5.1 g).

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.50 (d, 1H), 7.15 (d, 1H), 7.08 (d, 1H), 6.93 (m, 1H), 6.55 (q, H), 6.49 (s, 1H), 6.38 (m, 2H), 4.10 (broad, 2H), 3.81 (m, 1H), 3.65 (m, 1H), 3.24 (d, 1H), 3.01 (d, 1H), 2.64 (m, 2H), 2.23 (m, 2H), 2.00 (m, 3H), 1.75 (m, 4H), 1.46-1.27 (m, 3H)

#### Step 3.

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-methoxyacetylaminophenyl)piperidin-3-ol

30

The solution of 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-

aminophenyl)piperidin-3-ol (1.0 g, 2.24 mmol) obtained in Step 2 in 30 ml of tetrahydrofuran was cooled to 0°C. Sodium carbonate (350 mg, 3.36 mmol) and methoxyacetyl chloride (0.53 mg, 5.82 mmol) was added to the reaction mixture, which was then stirred for 4 hours. 20 ml of ice water was added to the reaction mixture, which was then concentrated and extracted with 50 ml of ethyl acetate. The resulting organic layer was dried and concentrated on sodium sulfate. The resulting residue was crystallized with ethyl acetate to give 1.1 g of the titled compound.

Step 4.

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-methoxyacetylaminophenyl)piperidin-3-one

To the solution of 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-methoxyacetylaminophenyl)piperidin-3-ol (1.1 g) obtained in Step 3 in 5 ml of dimethyl sulfoxide, was added dropwise 1.2 ml of triethylamine. The reaction mixture was cooled to 0°C. Pyridine-SO<sub>3</sub> complex (0.9 g) was added to the reaction mixture, which was then stirred for about 4 hours at room temperature. The reaction mixture was diluted with 20 ml of ethyl acetate and washed with brine. The resulting organic layer was dried and concentrated on sodium sulfate. The resulting residue was purified with column chromatography to give 820 mg of the titled compound.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 9.42 (s, 1H), 8.25 (d, 1H), 7.58 (d, 1H), 7.49 (s, 1H), 7.11 (m, 2H), 6.79 (m, 1H), 6.54 (d, 1H), 6.44 (s, 1H), 4.69 (m, 1H), 4.00 (s, 2H), 3.63 (d, 1H), 3.45 (m, 4H), 3.17 (m, 2H), 2.78 (m, 1H), 2.31 (broad, 2H), 2.00 (m, 3H), 1.75 (m, 2H), 1.52 (m, 2H), 1.38 (m, 1H)

Example 88:

Preparation of

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-methoxycarbonylaminophenyl)piperidin-3-one



In accordance with the same procedure as in Example 87, except that methylchloroformate was used instead of methoxyacetyl chloride in Step 3 thereof, 671 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.91 (d, 1H), 7.71 (s, 1H), 7.50 (m, 2H), 7.16 (m, 2H), 6.77 (m, 1H), 6.55 (d, 1H), 6.45 (s, 1H), 4.66 (m, 1H), 3.78 (s, 3H), 3.53 (d, 1H), 3.43 (d, 1H), 3.19 (t, 1H), 3.07 (d, 1H), 2.78 (m, 1H), 2.26 (m, 2H), 2.00 (m, 3H), 1.75 (m, 2H), 1.48 (m, 2H), 1.21 (m, 1H)

Example 89:

Preparation of

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-cyclopropylcarbonylaminophenyl)piperidin-3-one

In accordance with the same procedure as in Example 87, except that cyclopropylcarbonyl chloride was used instead of methoxyacetyl chloride in Step 3 thereof, 670 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.51 (s, 1H), 8.19 (d, 1H), 7.62 (d, 1H), 7.50 (s, 1H), 7.15 (m, 2H), 6.74 (m, 1H), 6.55 (d, 1H), 6.44 (s, 1H), 4.65 (m, 1H), 3.58 (d, 1H), 3.48 (d, 1H), 3.23 (t, 1H), 3.07 (d, 1H), 2.80 (m, 1H), 2.31 (broad, 2H), 2.00 (m, 3H), 1.75 (m, 2H), 1.52 (m, 2H), 1.38 (m, 1H), 1.09 (q, 3H), 0.88 (q, 3H)

Example 90:

Preparation of

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-isobutyrylaminophenyl)piperidin-3-one

In accordance with the same procedure as in Example 87, except that isobutyryl chloride was used instead of methoxyacetyl chloride in Step 3 thereof, 522 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.36 (s, 1H), 8.23 (dd, 1H), 7.59 (d, 1H),

7.50 (s, 1H), 7.15 (m, 2H), 6.75 (m, 1H), 6.55 (d, 1H), 6.45 (s, 1H), 4.65 (m, 1H), 3.55 (d, 1H), 3.46 (d, 1H), 3.25 (t, 1H), 3.07 (d, 1H), 2.80 (m, 1H), 2.31 (broad, 2H), 2.00 (m, 3H), 1.75 (m, 2H), 1.52 (m, 2H), 1.38 (m, 1H), 1.21 (m, 6H)

5 Example 91:

Preparation of

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(4-chlorobutyryl)aminophenyl]piperidin-3-one

10 In accordance with the same procedure as in Example 87, except that chlorobutyryl chloride was used instead of methoxyacetyl chloride in Step 3 thereof, 750 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.42 (s, 1H), 8.17 (d, 1H), 7.547(s, 1H), 7.31 (s, 1H), 7.12 (m, 2H), 6.75 (m, 1H), 6.55 (d, 1H), 6.45 (s, 1H), 4.92 (m, 1H), 15 3.55 (d, 1H), 3.46 (d, 1H), 3.25 (t, 1H), 3.07 (d, 1H), 2.80-2.52 (m, 5H), 2.44 (m, 2H), 2.31 (m, 2H), 2.00 (m, 3H), 1.88-1.58 (m, 4H), 1.52-1.22 (m, 5H)

Example 92:

Preparation of

20 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(3,3-dimethylazetidin-2-one-1-yl)phenyl]piperidin-3-one

Step 1.

25 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(3-chloro-2,2-dimethylpropionyl)aminophenyl]piperidin-3-ol

In accordance with the same procedure as in Step 3 of Example 87, except that 3-chloropivaloyl chloride was used instead of methoxyacetyl chloride, 1.1 g of the titled compound was prepared.

30

Step 2.

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluor

o-2-(3,3-dimethylazetidin-2-one-1-yl)phenyl]piperidin-3-ol

The solution of 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexane carbonyl]amino]-1-[4-fluoro-2-(3-chloro-2,2-dimethylpropionyl)aminophenyl]piperidin-3-ol (1.1 g) obtained in Step 1 in 40 ml of dichloromethane was cooled to 0°C and was added sodium tert-butoxide (NaOtBu, 600 mg) thereto. The reaction mixture was stirred overnight at room temperature, diluted with dichloromethane, and washed with brine. The resulting organic layer was dried and concentrated on sodium sulfate. The resulting residue was purified with column chromatography to give 762 mg of the titled compound.

### Step 3.

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(3,3-dimethylazetidin-2-one-1-yl)phenyl]piperidin-3-one

15

To the solution of 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexane carbonyl]amino]-1-[4-fluoro-2-(3,3-dimethylazetidin-2-one-1-yl)phenyl]piperidin-3-ol (0.5 g) obtained in Step 2 in 10 ml of dimethylsulfoxide, was added 0.5 ml of triethylamine. The reaction mixture was cooled to 0°C. Pyridine-SO<sub>3</sub> complex (0.45 g) was added to the reaction mixture, which was then stirred for about 4 hours at room temperature. The reaction mixture was diluted with 20 ml of ethyl acetate and washed with brine. The resulting organic layer was dried and concentrated on sodium sulfate. The resulting residue was filtered and dried to give 340 mg of the titled compound.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.81 (d, 1H), 7.50 (s, 2H), 7.15 (m, 2H), 6.88 (m, 1H), 6.54 (q, 1H), 6.45 (s, 1H), 6.63 (m, 1H), 3.78 (d, 1H), 3.67 (d, 1H), 3.55 (d, 1H), 3.27 (t, 1H), 3.19 (d, 1H), 2.78 (m, 1H), 2.26 (t, 2H), 1.97 (t, 2H), 1.73 (m, 4H), 1.48 (m, 2H), 1.36-1.21 (m, 7H)

### 30 Example 93:

Preparation of

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-

(3-methyl-aziridin-2-one-1-yl)phenyl]piperidin-3-one

In accordance with the same procedure as in Example 92, except that 2-chloropropionyl chloride was used instead of 3-chloropivaloyl chloride in Step 1 thereof, 750 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 9.69 (s, 1H), 8.60 (d, 1H), 7.49 (s, 2H), 7.14 (d, 2H), 6.81 (m, 1H), 6.54 (s, 1H), 6.48 (s, 1H), 4.67 (m, 1H), 4.54 (m, 1H), 3.65 (d, 1H), 3.59 (m, 1H), 3.44 (m, 1H), 2.76 (m, 1H), 2.21 (m, 2H), 1.95 (m, 2H), 1.81 (s, 3H), 1.71 (m, 4H), 1.44 (m, 2H), 1.36 (m, 1H)

Example 94:

Preparation of

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(azetidin-2-one-1-yl)phenyl]piperidin-3-one

In accordance with the same procedure as in Example 92, except that 3-chloropropionyl chloride was used instead of 3-chloropivaloyl chloride in Step 1 thereof, 750 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.44 (d, 1H), 8.32 (d, 1H), 7.52 (s, 2H), 7.14 (d, 2H), 6.81 (m, 1H), 6.54 (s, 1H), 6.48 (s, 1H), 4.60 (m, 1H), 3.65 (d, 1H), 3.59 (m, 1H), 3.44 (m, 1H), 3.00 (m, 2H), 2.76 (m, 1H), 2.21 (m, 2H), 2.01 (m, 2H), 1.95 (m, 2H), 1.71 (m, 4H), 1.44 (m, 2H), 1.36 (m, 1H)

Example 95:

Preparation of

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(pyrrolidin-2-one-1-yl)phenyl]piperidin-3-one

In accordance with the same procedure as in Example 92, except that 4-chlorobutyryl chloride was used instead of 3-chloropivaloyl chloride in Step 1 thereof, 230 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.49 (m, 1H), 7.26 (m, 2H), 7.03 (m, 3H),

6.53 (m, 1H) , 6.43 (s, 1H), 4.64 (m, 1H), 3.80 (m, 1H), 3.70 (m, 1H), 3.55 (d, 1H), 3.47 (d, 1H), 3.23 (m, 2H), 2.57 (m, 1H), 2.55 (d, 2H), 2.26 (m, 2H), 2.15 (m, 2H), 1.96 (m, 3H), 1.59 (m, 2H), 1.48 (m, 1H)

5 Example 96:

Preparation of

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(1,3-imidazolidin-2,5-dione-1-yl)phenyl]piperidin-3-one

10 Step 1.

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(ethoxycarbonylmethylcarbamoyl)aminophenyl]piperidin-3-ol

15 In accordance with the same procedure as in Step 3 of Example 87, except that ethyl isocyanatoacetate was used instead of methoxyacetyl chloride, 400 mg of the titled compound was prepared.

Step 2.

20 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(1,3-imidazolidin-2,5-dione-1-yl)aminophenyl]piperidin-3-ol

25 To the solution of 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(ethoxycarbonylmethylcarbamoyl)aminophenyl]piperidin-3-ol (1.1 g) obtained in Step 1 in 10 ml of ethanol, was added 5 ml of hydrochloride solution. The reaction mixture was refluxed for about 6 hours, diluted with ethyl acetate (20 ml), and washed with brine. The resulting organic layer was dried and concentrated on sodium sulfate. The resulting residue was filtered and dried to give 140 mg of the titled compound as white color.

30

Step 3.

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-

(1,3-imidazolidin-2,5-dione-1-yl)aminophenyl]piperidin-3-one

To the solution of 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexane carbonyl]amino]-1-[4-fluoro-2-(1,3-imidazolidin-2,5-dione-1-yl)aminophenyl]piperidin-3-ol (140 mg) obtained in Step 2 in 10 ml of dimethylsulfoxide, was added 0.2 ml of triethylamine. The reaction mixture was cooled to 0°C. Pyridine-SO<sub>3</sub> complex (0.2 g) was added slowly to the reaction mixture, which was then stirred for about 4 hours at room temperature. The reaction mixture was diluted with 20 ml of ethyl acetate and washed with brine. The resulting organic layer was dried and concentrated on sodium sulfate. The resulting residue was filtered and dried to give 83 mg of the titled compound as white color.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.96 (d, 1H), 7.85 (s, 1H), 7.66 (m, 1H), 7.55 (m, 1H), 7.53 (s, 2H), 7.13 (d, 1H), 6.99 (s, 1H), 6.61 (m, 1H), 6.59 (s, 1H), 6.43 (s, 1H), 4.78 (m, 1H), 4.26 (d, 2H), 4.01 (m, 2H), 3.83 (m, 2H), 3.11 (d, 1H), 2.98 (m, 1H), 2.76 (m, 1H), 2.21 (m, 2H), 1.95 (m, 2H), 1.71 (m, 4H), 1.44 (m, 2H), 1.36 (m, 1H)

Example 97:

Preparation of

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(1,1-dioxo-isothiazolidin-1-yl)phenyl]piperidin-3-one

In accordance with the same procedure as in Example 92, except that 3-chloropropanesulfonyl chloride was used instead of 3-chloropivaloyl chloride in Step 1 thereof, 80 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.27 (s, 1H), 8.18 (d, 1H), 7.62 (d, 1H), 7.50 (s, 1H), 7.15 (m, 2H), 6.76 (m, 1H), 6.55 (d, 1H), 6.46 (s, 1H), 4.62 (m, 1H), 3.56 (d, 1H), 3.45 (d, 1H), 3.23 (t, 1H), 3.00 (m, 1H), 2.78 (m, 1H), 2.27 (m, 4H), 1.98 (m, 2H), 1.73 (m, 6H), 1.48 (m, 2H), 1.38 (m, 1H)

Example 98:

## Preparation of

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(piperidin-2-one-1-yl)phenyl]piperidin-3-one

5 In accordance with the same procedure as in Example 92, except that 5-chlorovaleryl chloride was used instead of 3-chloropivaloyl chloride in Step 1 thereof, 130 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.50 (s, 2H), 7.29 (m, 2H), 7.00 (m, 1H), 6.92 (d, 1H), 6.54 (d, 1H), 6.47 (s, 1H), 4.63 (m, 1H), 3.67 (m, 2H), 3.52-3.27 (m, 4H), 2.96 (s, 1H), 2.88 (s, 1H), 2.55 (m, 2H), 2.26 (m, 2H), 2.00 (m, 2H), 1.98-1.75 (m, 6H), 1.66 (m, 4H), 1.44-1.25 (m, 4H)

Example 99:

## Preparation of

15 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(pyrrol-1-yl)phenyl]piperidin-3-one

## Step 1.

20 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(pyrrol-1-yl)phenyl]piperidin-3-ol

The mixture of 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-aminophenyl)piperidin-3-ol (3.0 g) obtained in Step 2 of Example 87 and 2,5-dimethoxytetrahydrofuran (0.96 ml) was refluxed in 80 ml of acetic acid for 1 hour. The reaction mixture was concentrated under reduced pressure. The resulting residue was diluted with 70 ml of dichloromethane and washed with a saturated sodium bicarbonate and brine. The resulting organic layer was dried and concentrated on sodium sulfate. The resulting residue was purified with column chromatography to give 1.8 g of the titled compound as white color.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.50 (s, 2H), 7.29 (m, 2H), 7.00 (m, 1H), 6.92 (d, 1H), 6.54 (d, 1H), 6.47 (s, 1H), 6.26 (s, 2H), 4.00 (broad, 1H), 3.71 (m,

2H), 3.51 (m, 1H), 3.19 (d, 1H), 2.78 (s, 1H), 2.55 (m, 2H), 2.26 (m, 2H), 2.00 (m, 2H), 1.98-1.75 (m, 6H), 1.66 (m, 4H), 1.44-1.25 (m, 4H)

Step 2.

- 5 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(pyrrol-1-yl)phenyl]piperidin-3-one

To the solution of 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexane  
carbonyl]amino]-1-[4-fluoro-2-(pyrrol-1-yl)phenyl]piperidin-3-ol (1.8 g) obtained  
10 in Step 1 in 10 ml of dimethylsulfoxide, was added 2 ml of triethylamine. The  
reaction mixture was cooled to 0°C. Pyridine-SO<sub>3</sub> complex (1.8 g) was added  
to the reaction mixture, which was then stirred for about 2 hours at room  
temperature. The reaction mixture was diluted with 50 ml of ethyl acetate and  
washed with brine. The resulting organic layer was dried and concentrated on  
15 sodium sulfate. The resulting residue was filtered and dried to give 1.5 g of  
the titled compound as white color.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.48 (s, 1H), 7.38 (d, 1H), 7.14 (d, 1H),  
6.95 (m, 5H), 6.41 (d, 1H), 6.38 (s, 1H), 6.29 (q, 2H), 4.57 (m, 1H), 3.43 (s, 2H),  
2.93-2.85 (m, 2H), 2.51 (m, 1H), 2.23 (d, 2H), 1.95 (t, 2H), 1.76-1.53 (m, 4H),  
20 1.52-1.22 (m, 3H)

Example 100:

Preparation of

- 25 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(oxazol-4-yl)phenyl]piperidin-3-one

Step 1.

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(oxazol-4-yl)phenyl]piperidin-3-ol

30

The mixture of  
4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-



formylphenyl)piperidin-3-ol (3.0 g, 6 mmol) obtained in Example 79, potassium carbonate (2 g, 20 mmol) and p-toluenesulfonylmethyl isocyanate (TosMIC; 1.8 g, 9 mmol) was refluxed in 500 ml of methanol for about 2 hours. The reaction mixture was concentrated, added water thereto, and extracted with dichloromethane. The resulting organic layer was dried and concentrated on sodium sulfate. The resulting residue was purified with column chromatography to give 3.2 g of the titled compound.

Step 2.

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(oxazol-4-yl)phenyl]piperidin-3-ol

To the solution of 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(oxazol-4-yl)phenyl]piperidin-3-ol (1.8 g) obtained in Step 1 in 20 ml of dimethylsulfoxide, was added 2.7 ml of triethylamine (19 mmol). The reaction mixture was cooled to 0°C. Pyridine-SO<sub>3</sub> complex (3.04 g, 19 mmol) was added to the reaction mixture, which was then stirred for about 2 hours at room temperature. The reaction mixture was diluted with 100 ml of ethyl acetate and washed with brine. The resulting organic layer was dried and concentrated on sodium sulfate. The resulting residue was purified with column chromatography to give 1.67 g of the titled compound.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.91 (s, 1H), 7.65 (s, 1H), 7.48 (m, 3H), 7.22 (m, 2H), 7.00 (m, 1H), 6.54 (d, 2H), 4.67 (m, 1H), 3.56 (s, 2H), 3.22 (m, 2H), 2.78 (m, 1H), 2.29 (d, 2H), 1.98 (m, 2H), 1.84-1.62 (m, 5H), 1.43-1.19 (m, 3H)

Example 101:

Preparation of

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(N-methylcarbamoyl)phenyl]piperidin-3-one

## Step 1.

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(N-methylcarbamoyl)phenyl]piperidin-3-ol

5

To the solution of 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-formylphenyl)piperidin-3-ol (4.0 g, 6 mmol) obtained in Example 79 and potassium bicarbonate ( $\text{NaH}_2\text{PO}_4$ ; 340 mg) in the mixed solvent of 20 ml of acetonitrile and 6 ml of distilled water, was 35%-hydrogen peroxide solution (1.0 ml) and sodium hypochlorite ( $\text{NaClO}_4$ ; 1.2 g) at 10°C. The reaction mixture was stirred for about 1 hour at room temperature, added sodium sulfite (1.0 g) thereto, and acidified with 10%-hydrochloride solution. The resulting solid was filtered and dried to give 3.6 g of a solid. The resulting solid (2.5 g), monomethylamine hydrochloride (1.0 g) and benzotriazole-1-yloxytripyrrolidinophosphonium hexaphosphate (PyBOP; 800 mg) were dissolved in 30 ml of dichloromethane (MC; 30 ml), cooled to 0°C. Diisopropylethylamine (DIEA; 1.0 g) was added to the reaction mixture, which was then stirred for about 4 hours. Water was added to the reaction mixture. The resulting organic layer was dried and concentrated on sodium sulfate. The resulting residue was purified with column chromatography to give 1.5 g of the titled compound.

10  
15  
20

$^1\text{H-NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  10.25 (broad, 1H), 7.89 (d, 1H), 7.51 (s, 1H), 7.22-7.16 (m, 4H), 6.56 (m, 2H), 4.15 (m, 2H), 3.91-3.79 (m, 2H), 3.35 (d, 1H), 3.21 (d, 1H), 2.98 (s, 3H), 2.76 (m, 2H), 2.23 (m, 2H), 2.00 (m, 4H), 1.70 (m, 4H), 1.40-1.20 (m, 3H)

25

## Step 2.

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(N-methylcarbamoyl)phenyl]piperidin-3-ol

30

To the solution of 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexane

carbonyl]amino]-1-[4-fluoro-2-(N-methylcarbamoyl)phenyl]piperidin-3-ol (1.8 g) obtained in Step 1 in 10 ml of dimethyl sulfoxide, was added 2 ml of triethylamine. The reaction mixture was cooled to 0°C. Pyridine-SO<sub>3</sub> complex (1.8 g) was added slowly to the reaction mixture, which was then stirred for  
5 about 2 hours at room temperature. The reaction mixture was diluted with 50 ml of ethyl acetate and washed with brine. The resulting organic layer was dried and concentrated on sodium sulfate. The resulting residue was purified with column chromatography to give 1.2 g of the titled compound.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.86 (d, 1H), 7.85 (d, 1H), 7.62 (d, 1H),  
10 7.50 (s, 1H), 7.16 (m, 3H), 6.55 (d, 1H), 6.46 (s, 1H), 4.56 (d, 1H), 3.62 (s, 2H),  
3.26 (d, 2H), 2.97 (s, 3H), 2.78 (m, 1H), 2.68 (s, 1H), 2.27(m, 2H), 1.70 (t, 3H),  
1.43-1.21 (m, 3H)

#### Example 102:

15 Preparation of  
4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(N,N-di  
methylcarbamoyl)phenyl]piperidin-3-one

In accordance with the same procedure as in Example 101, except that  
20 dimethylamine was used instead of monomethylamine hydrochloride in Step 1 thereof, 1.05 g of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.49 (m, 2H), 7.13 (s, 1H), 6.99 (m, 3H),  
6.54 (q, 1H), 6.43 (s, 1H), 4.61 (m, 1H), 3.73 (m, 1H),, 3.53 (m, 2H), 3.07 (s,  
3H), 2.75 (s, 3H), 2.58 (broad, 1H), 2.24 (m, 2H), 1.96 (m, 2H), 1.72 (m, 4H),  
25 1.47-1.24 (m, 3H)

#### Example 103:

Preparation of  
4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(N-cyclo  
30 propylcarbamoyl)phenyl]piperidin-3-one

In accordance with the same procedure as in Example 101, except that

cyclopropylamine was used instead of monomethylamine hydrochloride in Step 1 thereof, 240 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 9.11 (s, 1H), 7.88 (d, 1H), 7.62 (d, 1H), 7.50 (d, 1H), 7.13 (m, 3H), 6.55 (d, 1H), 6.47 (s, 1H), 4.67 (m, 1H), 3.58 (m, 2H), 3.26 (m, 2H), 2.99 (m, 1H), 2.87 (m, 1H), 2.78 (m, 1H), 2.25 (m, 2H), 1.99 (m, 2H), 1.50 (m, 4H), 1.42-1.23 (m, 3H), 0.88 (m, 2H), 0.55 (m, 2H)

Example 104:

Preparation of

4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-aminophenyl)piperidin-3-one

Step 1.

4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-aminophenyl)piperidin-3-ol

In accordance with the same procedure as in Step 2 of Example 87, except that

4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-nitrophenyl)piperidin-3-ol was used instead of 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-nitrophenyl)piperidin-3-ol, 1.1 g of the titled compound was prepared.

Step 2.

4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-aminophenyl)piperidin-3-one

To the solution of 4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-aminophenyl)piperidin-3-ol (1.1 g) obtained in Step 1 in 5 ml of dimethyl sulfoxide, was added 1.2 ml of triethylamine. The reaction mixture was cooled to 0°C. Pyridine-SO<sub>3</sub> complex (0.9 g) was added to the reaction mixture, which was then stirred for about 4 hours at room

temperature. The reaction mixture was diluted with 20 ml of ethyl acetate and washed with brine. The resulting organic layer was dried and concentrated on sodium sulfate. The resulting residue was filtered and dried to give 820 mg of the titled compound as white color.

5           <sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.11 (m, 1H), 6.88 (m, 1H), 6.56 (d, 1H), 6.48 (s, 1H), 6.34 (m, 1H), 6.21 (s, 1H), 4.59 (s, 1H), 4.22 (m, 1H), 3.16 (m, 4H), 2.80 (d, 1H), 2.52 (s, 3H), 2.23 (d, 1H), 2.00-1.78 (m, 4H), 1.68 (m, 2H), 1.42-1.21 (m, 3H)

10    Example 105:

Preparation of

4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-nitrophenyl)piperidin-3-one

15           In accordance with the same procedure as in Example 76, except that 1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, and 1-fluoronitrobenzene was used instead of 1-fluorophenyl methyl sulfone in Step 5 thereof, 330 mg of the titled compound was prepared.

20           <sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.10 (d, 1H), 7.66 (t, 1H), 7.42 (m, 2H), 7.32 (d, 1H), 6.93 (s, 1H), 6.44 (s, 1H), 4.70 (m, 1H), 3.86, (q, 3H), 3.41 (t, 1H), 2.78 (m, 1H), 2.50 (s, 3H), 2.31 (broad, 2H), 2.00 (m, 3H), 1.75 (m, 2H), 1.52 (m, 2H), 1.38 (m, 1H)

25    Example 106:

Preparation of

4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

30           In accordance with the same procedure as in Example 76, except that 1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in

Step 3 thereof, 870 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.10 (d, 1H), 7.66 (t, 1H), 7.42 (m, 2H),  
7.32 (d, 1H), 6.93 (s, 1H), 6.44 (s, 1H), 4.71 (m, 1H), 3.78 (d, 1H), 3.65 (d, 1H),  
3.51 (d, 1H), 3.18 (s, 3H), 2.69 (m, 1H), 2.50 (s, 3H), 2.20 (t, 2H), 1.98 (m, 3H),  
5 1.60 (m, 2H), 1.47 (m, 2H), 1.38(m, 1H)

Example 107:

Preparation of

4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1  
10 -(2-ethanesulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that  
1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexanecarboxylic acid was  
used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in  
15 Step 3 thereof, and 1-fluorophenyl ethyl sulfone was used instead of  
1-fluorophenyl methyl sulfone in Step 5 thereof, 330 mg of the titled compound  
was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.07 (d, 1H), 7.65 (t, 1H), 7.39 (m, 2H),  
7.31 (d, 1H), 6.93 (s, 1H), 6.44 (s, 1H), 4.70 (m, 1H), 3.77 (d, 1H), 3.61 (d, 1H),  
20 3.57 (d, 1H), 3.39 (m, 2H), 3.11 (t, 1H), 2.78 (m, 1H), 2.50 (s, 3H), 2.24 (t, 2H),  
2.00 (t, 2H), 1.89 (m, 1H), 1.70 (m, 2H), 1.47 (m, 2H), 1.31 (m, 1H), 1.10 (t, 3H)

Example 108:

Preparation of

25 4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1  
-[4-fluoro-2-(3,3-dimethyl-azetidin-2-one-1-yl)phenyl]piperidin-3-one

In accordance with the same procedure as in Example 92, except that  
4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1  
30 -[4-fluoro-2-(3-chloro-2,2-dimethylpropionyl)aminophenyl]piperidin-3-ol was  
used instead of  
4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-

(3-chloro-2,2-dimethylpropionyl)aminophenyl]piperidin-3-ol in Step 2 thereof, 240 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.80 (d, 1H), 7.34 (d, 1H), 7.10 (m, 1H), 6.95 (s, 1H), 6.80 (m, 1H), 6.43 (s, 1H), 4.63 (m, 1H), 3.67 (d, 1H), 3.55 (m, 2H), 3.50 (d, 1H), 3.43 (t, 1H), 3.21 (d, 1H), 2.78 (m, 1H), 2.50 (s, 3H), 2.24 (t, 2H), 1.96 (t, 2H), 1.74-1.69 (m, 4H), 1.43-1.20 (m, 10H)

#### Example 109:

##### Preparation of

4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(pyrrolidin-2-one-1-yl)phenyl]piperidin-3-one

In accordance with the same procedure as in Example 92, except that 4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(4-chlorobutyryl)aminophenyl]piperidin-3-ol was used instead of 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(3-chloro-2,2-dimethylpropionyl)aminophenyl]piperidin-3-ol in Step 2 thereof, 130 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.34 (d, 1H), 7.09-6.90 (m, 4H), 6.43 (s, 1H), 4.62 (m, 1H), 3.81 (m, 1H), 3.70 (m, 1H), 3.55 (q, 2H), 3.23 (m, 2H), 2.78 (m, 1H), 2.62-2.50 (m, 7H), 2.23 (m, 5H), 1.94 (t, 2H), 1.70-1.58 (m, 4H), 1.40-1.21 (m, 3H)

#### Example 110:

##### Preparation of

4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(oxazolidin-2-one-3-yl)phenyl]piperidin-3-one

In accordance with the same procedure as in Example 92, except that 4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(2-chloro-ethoxycarbonyl)aminophenyl]piperidin-3-ol was used instead of

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(3-chloro-2,2-dimethylpropionyl)aminophenyl]piperidin-3-ol in Step 2 thereof, 130 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.06 (s, 1H), 6.88 (q, 1H), 6.54 (d, 1H),  
5 6.47 (s, 1H), 6.32 (t, 1H), 6.20 (d, 1H), 5.62 (s, 1H), 4.53 (s, 1H), 4.22 (m, 1H),  
3.16 (m, 3H), 2.84 (d, 1H), 2.52 (s, 3H), 2.23 (d, 2H), 2.11-1.83 (m, 3H),  
1.64-1.52 (m, 4H), 1.48-1.27 (m, 6H)

Example 111:

10 Preparation of

4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(imidazolidin-2-one-1-yl)phenyl]piperidin-3-one

In accordance with the same procedure as in Example 92, except that  
15 4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(2-chloroethylcarbamoylamino)phenyl]piperidin-3-ol was used instead of  
4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(3-chloro-2,2-dimethylpropionyl)aminophenyl]piperidin-3-ol in Step 2 thereof, 79  
20 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.14 (d, 1H), 7.03-6.91 (m, 3H), 6.43 (s, 1H),  
4.69 (m, 1H), 3.93-3.51 (m, 8H), 3.43 (m, 1H), 3.21 (t, 1H), 2.78 (m, 1H),  
2.50 (s, 3H), 2.23 (m, 2H), 1.96 (m, 2H), 1.73 (m, 2H), 1.46-1.26 (m, 3H)

25 Example 112:

Preparation of

4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(3,3-dimethyl-azetidin-2-one-1-yl)phenyl]piperidin-3-one

30 In accordance with the same procedure as in Example 92, except that  
4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(3-chloro-2,2-dimethylpropionyl)aminophenyl]piperidin-3-ol was used instead



of

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(3-chloro-2,2-dimethylpropionyl)aminophenyl]piperidin-3-ol in Step 2 thereof, 373 mg of the titled compound was prepared.

5  $^1\text{H-NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  9.62 (s, 1H), 7.87 (f, 1H), 7.11 (q, 3H), 6.87 (m, 2H), 6.41 (s, 1H), 4.98 (m, 1H), 4.11 (m, 2H), 3.69 (m, 2H), 3.31 (d, 1H), 3.14 (d, 1H), 2.78 (m, 1H), 2.50 (s, 3H), 2.17 (m, 3H), 1.88 (q, 2H), 1.67 (m, 5H), 1.45-1.26 (m, 12H)

10 Example 113:

Preparation of

4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(pyrrolidin-2-one-1-yl)phenyl]piperidin-3-one

15 In accordance with the same procedure as in Example 92, except that 4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(4-chlorobutyrylamino)phenyl]piperidin-3-ol was used instead of 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(3-chloro-2,2-dimethylpropionyl)aminophenyl]piperidin-3-ol in Step 2 thereof, 20 373 mg of the titled compound was prepared.

$^1\text{H-NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  7.33 (m, 3H), 7.15 (m, 2H), 6.94 (s, 1H), 6.43 (s, 1H), 4.65 (m, 1H), 3.80 (m, 1H), 3.66 (m, 2H), 3.55 (d, 1H), 3.41 (d, 1H), 3.23 (t, 1H), 2.78 (m, 1H), 2.54 (m, 2H), 2.50 (s, 1H), 2.23 (m, 6H), 1.96 (t, 2H), 1.66 (m, 4H), 1.45-1.21 (m, 3H)

25

Example 114:

Preparation of

4-[N-[1-[N-(5-chloro-thiophen-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

30

In accordance with the same procedure as in Example 76, except that 1-[N-(5-chloro-thiophen-2-ylcarbonyl)amino]cyclohexanecarboxylic acid was

used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 1.1 g of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.08 (d, 1H), 7.65 (t, 1H), 7.45 (m, 3H), 7.32 (d, 1H), 6.92 (d, 1H), 5.95 (s, 1H), 4.67 (m, 1H), 3.77 (d, 1H), 3.65 (d, 1H), 3.55 (m, 1H), 3.19 (m, 1H), 3.18 (s, 3H), 2.69 (m, 1H), 2.24 (d, 2H), 2.00 (m, 3H), 1.65 (m, 3H), 1.45 (m, 3H)

#### Example 115:

Preparation of

4-[N-[1-[N-(5-chloro-thiophen-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-ethanesulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that 1-[N-(5-chloro-thiophen-2-ylcarbonyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, and 1-fluorophenyl ethyl sulfone was used instead of 1-fluorophenyl methyl sulfone in Step 5 thereof, 450 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.06 (d, 1H), 7.65 (t, 1H), 7.45 (m, 4H), 6.92 (d, 1H), 5.96 (d, 1H), 4.66 (m, 1H), 3.77 (d, 1H), 3.63 (d, 1H), 3.53 (m, 1H), 3.32 (m, 2H), 3.17 (m, 1H), 2.69 (m, 1H), 2.23 (d, 2H), 1.97 (m, 3H), 1.65 (m, 3H), 1.45 (m, 3H), 1.12 (t, 3H)

#### Example 116:

Preparation of

4-[N-[1-[N-(5-chloro-thiophen-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-isopropanesulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that 1-[N-(5-chloro-thiophen-2-ylcarbonyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, and 1-fluorophenyl isopropane sulfone was used instead of

1-fluorophenyl methyl sulfone in Step 5 thereof, 560 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.05 (d, 1H), 7.65 (t, 1H), 7.45 (m, 3H), 7.32 (d, 1H), 6.92 (d, 1H), 5.99 (s, 1H), 4.66 (m, 1H), 3.79 (m, 1H), 3.77 (d, 1H), 3.59 (d, 1H), 3.53 (m, 1H), 3.14 (m, 1H), 2.69 (m, 1H), 2.24 (d, 2H), 1.89 (m, 3H), 1.70 (m, 3H), 1.43 (m, 3H), 1.30 (d, 3H), 1.08 (d, 3H)

Example 117:

Preparation of

4-[N-[1-[N-(5-chloro-thiophen-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(5-fluoro-2-methanesulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that 1-[N-(5-chloro-thiophen-2-ylcarbonyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, and 2,4-difluorophenyl methane sulfone was used instead of 1-fluorophenyl methyl sulfone in Step 5 thereof, 210 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.80 (m, 1H), 7.35 (m, 5H), 6.93 (d, 1H), 5.93 (s, 1H), 4.66 (m, 1H), 3.77 (d, 1H), 3.65 (d, 1H), 3.52 (m, 1H), 3.19 (s, 3H), 3.18 (m, 1H), 2.69 (m, 1H), 2.24 (d, 2H), 2.00 (m, 3H), 1.70 (m, 3H), 1.42 (m, 3H)

Example 118:

Preparation of

4-[N-[1-[N-(5-methyl-thiophen-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that 1-[N-(5-methyl-thiophen-2-ylcarbonyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 562 mg of the titled compound was prepared.

$^1\text{H-NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  8.08 (d, 1H), 7.60 (m, 2H), 7.40 (m, 4H), 6.76 (d, 1H), 6.01 (s, 1H), 4.68 (m, 1H), 3.77 (d, 1H), 3.65 (d, 1H), 3.55 (m, 1H), 3.19 (m, 1H), 3.18 (s, 3H), 2.69 (m, 1H), 2.52 (s, 3H), 2.24 (d, 2H), 1.95 (m, 3H), 1.70 (m, 3H), 1.45 (m, 3H)

5

Example 119:

Preparation of

4-[N-[1-[N-(5-methyl-thiophen-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-isopropanesulfonylphenyl)piperidin-3-one

10

In accordance with the same procedure as in Example 76, except that 1-[N-(5-methyl-thiophen-2-ylcarbonyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, and 1-fluorophenyl isopropane sulfone was used instead of 1-fluorophenyl methyl sulfone in Step 5 thereof, 110 mg of the titled compound was prepared.

15

$^1\text{H-NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  8.21 (m, 1H), 7.55 (m, 2H), 7.33 (m, 3H), 6.79 (d, 1H), 5.92 (s, 1H), 4.66 (m, 1H), 3.79 (m, 1H), 3.77 (d, 1H), 3.59 (d, 1H), 3.53 (m, 1H), 3.17 (m, 1H), 2.69 (m, 1H), 2.53 (s, 3H), 2.24 (m, 2H), 1.89 (m, 3H), 1.70 (m, 3H), 1.43 (m, 3H), 1.30 (d, 3H), 1.08 (d, 3H)

20

Example 120:

Preparation of

4-[N-[1-[N-(5-methyl-thiophen-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(5-fluoro-2-methanesulfonylphenyl)piperidin-3-one

25

In accordance with the same procedure as in Example 76, except that 1-[N-(5-methyl-thiophen-2-ylcarbonyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, and 2,4-difluorophenyl methane sulfone was used instead of 1-fluorophenyl methyl sulfone in Step 5 thereof, 222 mg of the titled compound was prepared.

30

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.55 (m, 1H), 7.45 (m, 1H), 7.35 (m, 3H), 7.15 (m, 1H), 6.75 (d, 1H), 5.95 (s, 1H), 4.68 (m, 1H), 3.75 (d, 1H), 3.63 (d, 1H), 3.52 (m, 1H), 3.19 (s, 3H), 3.18 (m, 1H), 2.69 (m, 1H), 2.52 (s, 3H), 2.24 (d, 2H), 1.95 (m, 3H), 1.70 (m, 3H), 1.42 (m, 3H)

5

Example 121:

Preparation of

4-[N-[1-[N-(3-methyl-thiophen-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

10

In accordance with the same procedure as in Example 76, except that 1-[N-(3-methyl-thiophen-2-ylcarbonyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 131 mg of the titled compound was prepared.

15

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.09 (d, 1H), 7.66 (m, 1H), 7.54 (d, 1H), 7.42 (m, 2H), 7.31 (d, 1H), 6.91 (d, 1H), 5.95 (s, 1H), 4.70 (m, 1H), 3.80 (d, 1H), 3.63 (d, 1H), 3.41 (m, 1H), 3.29 (td, 1H), 3.18 (s, 3H), 2.78 (m, 1H), 2.55 (s, 3H), 2.27 (m, 2H), 1.95-1.82 (m, 3H), 1.72 (m, 2H), 1.48 (m, 2H), 1.32 (m, 1H)

20

Example 122:

Preparation of

4-[N-[1-[N-(2-methyl-furan-3-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

25

In accordance with the same procedure as in Example 76, except that 1-[N-(2-methyl-furan-3-ylcarbonyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 180 mg of the titled compound was prepared.

30

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.09 (d, 1H), 7.66 (q, 1H), 7.54 (d, 1H), 7.42 (m, 2H), 7.31 (d, 1H), 6.92 (d, 1H), 5.94 (s, 1H), 4.70 (m, 1H), 3.80 (d, 1H), 3.63 (d, 1H), 3.41 (m, 1H), 3.29 (td, 1H), 3.18 (s, 3H), 2.78 (m, 1H), 2.55 (s, 3H), 2.27 (m, 2H), 1.95-1.82 (m, 3H), 1.72 (m, 2H), 1.48 (m, 2H), 1.32 (m, 1H)

Example 123:

Preparation of

4-[N-[1-[N-(5-methyl-furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2  
5 -methanesulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that  
1-[N-(5-methyl-furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid was used  
instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3  
10 thereof, 123 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.09 (d, 1H), 7.64 (m, 1H), 7.54 (d, 1H),  
7.42 (m, 2H), 7.04 (d, 1H), 6.34 (s, 1H), 6.13 (d, 1H), 4.70 (m, 1H), 3.80 (d,  
1H), 3.63 (d, 1H), 3.41 (m, 1H), 3.29 (dt, 1H), 3.18 (s, 3H), 2.78 (m, 1H), 2.40 (s,  
3H), 2.27 (m, 2H), 1.95-1.82 (m, 3H), 1.72 (m, 2H), 1.48 (m, 2H), 1.32 (m, 1H)

15

Example 124:

Preparation of

4-[N-[1-[N-(5-bromo-furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2  
20 -methanesulfonyl-4-methoxyphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that  
1-[N-(5-methyl-furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid was used  
instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3  
25 thereof, and 2-fluoro-5-methoxyphenyl methane sulfone was used instead of  
1-fluorophenyl methyl sulfone in Step 5 thereof, 254 mg of the titled compound  
was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.8 (m, 1H), 7.48 (m, 1H), 7.38 (m, 2H),  
7.10 (s, 1H), 6.48 (s, 1H), 6.38 (s, 1H), 4.63 (s, 1H), 3.21 (s, 3H), 2.18-1.31 (m,  
30 12H)

Example 125:

## Preparation of

4-[N-[1-[N-(4,5-dimethyl-furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

5 In accordance with the same procedure as in Example 76, except that 1-[N-(4,5-dimethyl-furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 270 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.09 (d, 1H), 7.66 (m, 1H), 7.54 (d, 1H),  
10 7.42 (m, 2H), 6.92 (s, 1H), 6.32 (s, 1H), 4.70 (m, 1H), 3.80 (d, 1H), 3.63 (d, 1H), 3.41 (m, 1H), 3.29 (td, 1H), 3.18 (s, 3H), 2.78 (m, 1H), 2.55 (s, 3H), 2.29 (s, 3H), 2.27 (m, 2H), 2.01 (s, 3H), 1.95-1.82 (m, 3H), 1.72 (m, 2H), 1.48 (m, 2H), 1.32 (m, 1H)

15 Example 126:

## Preparation of

4-[N-[1-[N-(4,5-dimethyl-furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(pyrrolidin-2-one-1-yl)phenyl]piperidin-3-one

20 In accordance with the same procedure as in Example 92, except that 4-[N-[1-[N-(4,5-dimethyl-furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(4-chlorobutyryl)aminophenyl]piperidin-3-ol was used instead of 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(3-chloro-2,2-dimethylpropionyl)aminophenyl]piperidin-3-ol in Step 2 thereof,  
25 440 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.23 (m, 2H), 7.09 (m, 2H), 6.91 (s, 1H),  
4.65 (m, 1H), 3.71 (m, 1H), 3.62 (m, 2H), 3.51 (m, 2H), 3.33 (d, 1H), 3.21 (t, 1H),  
2.69 (m, 1H), 2.54 (m, 2H), 2.28 (d, 3H), 2.22 (m, 2H), 1.95 (s, 3H), 1.75 (m,  
4H), 1.43-1.21 (m, 3H)

30

Example 127:

## Preparation of

4-[N-[1-[N-(morpholin-4-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that 1-[N-(morpholin-4-ylcarbonyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 80 mg of the titled compound was prepared.

$^1\text{H-NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  8.1(m,1H), 7.6(m, 1H), 7.3(m, 2H), 6.8(d, 1H), 6.4(s, 1H), 4.7(m, 1H), 3.7(dd, 2H), 3.1(s, 3H), 1.4-2.0(m, 10H)

Example 128:

Preparation of

4-[N-[1-[N-(benzothiophen-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that 1-[N-(benzothiophen-2-ylcarbonyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 250 mg of the titled compound was prepared.

$^1\text{H-NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  8.07 (d, 1H), 7.87 (m, 3H), 7.65 (m, 1H), 7.45-7.39 (m, 5H), 6.25 (s, 1H), 4.70 (m, 1H), 3.80 (d, 1H), 3.63 (d, 1H), 3.41 (m, 1H), 3.29 (td, 1H), 3.18 (s, 3H), 2.78 (m, 1H), 2.27 (m, 2H), 1.95-1.82 (m, 3H), 1.72 (m, 2H), 1.48 (m, 2H), 1.32 (m, 1H),

Example 129:

Preparation of

4-[N-[1-[N-(benzothiophen-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-tert-butoxycarbonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 85, except that 4-[N-[1-[N-(benzothiophen-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-3-piperidinol hydrochloride was used instead of



4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-3-piperidinol hydrochloride and 2-tert-butyl bromobenzoate was used instead of 1-bromo-4-fluoro-2-thiomethoxybenzene in Step 1 thereof, 110 mg of the titled compound was prepared.

5  $^1\text{H-NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  7.84 (m, 2H), 7.74 (s, 1H), 7.51 (d, 1H), 7.49 (d, 1H), 7.43 (f, 1H), 7.32 (dd, 1H), 6.94 (m, 2H), 6.23 (s, 1H), 4.66 (m, 1H), 3.80 (d, 1H), 3.63 (d, 1H), 3.41 (m, 1H), 3.29 (td, 1H), 2.78 (m, 1H), 2.27 (m, 2H), 1.95-1.82 (m, 3H), 1.72 (m, 2H), 1.55 (s, 9H), 1.48 (m, 2H), 1.32 (m, 1H)

10

Example 130:

Preparation of

4-[N-[1-[N-(benzothiophen-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-chloro-5-trifluoromethylphenyl)piperidin-3-one

15

In accordance with the same procedure as in Example 85, except that 4-[N-[1-[N-(benzothiophen-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-3-piperidinol hydrochloride was used instead of 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-3-piperidinol hydrochloride and 1-bromo-2-chloro-5-trifluoromethylbenzene was used instead of 1-bromo-4-fluoro-2-thiomethoxybenzene in Step 1 thereof, 222 mg of the titled compound was prepared

20  $^1\text{H-NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  7.86 (m, 3H), 7.46 (m, 3H), 7.23 (m, 2H), 6.20 (s, 1H), 4.72 (m, 1H), 3.83 (d, 1H), 3.79 (d, 1H), 3.62 (d, 1H), 3.25 (t, 1H), 2.78 (m, 1H), 2.27 (m, 2H), 2.01 (t, 2H), 1.82-1.61 (m, 4H), 1.20 (m, 2H), 1.18 (m, 1H)

25

Example 131:

Preparation of

30 4-[N-[1-[N-(benzothiophen-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-chloro-2-fluoro-phenyl)piperidin-3-one

In accordance with the same procedure as in Example 85, except that 4-[N-[1-[N-(benzothiophen-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-3-piperidinol hydrochloride was used instead of 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-3-piperidinol hydrochloride and 1-bromo-2-fluoro-4-chlorobenzene was used instead of 1-bromo-4-fluoro-2-thiomethoxybenzene in Step 1 thereof, 76 mg of the titled compound was prepared

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.86 (m, 3H), 7.42 (m, 3H), 7.06 (m, 2H), 6.83 (d, 1H), 6.24 (s, 1H), 4.67 (m, 1H), 3.82 (d, 1H), 3.78 (d, 1H), 3.62 (d, 1H), 3.27 (t, 1H), 2.78 (m, 1H), 2.27 (m, 2H), 2.01 (t, 2H), 1.82-1.61 (m, 4H), 1.20 (m, 2H), 1.18 (m, 1H)

#### Example 132:

Preparation of

4-[N-[1-[N-(benzothiophen-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 85, except that 4-[N-[1-[N-(benzothiophen-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-3-piperidinol hydrochloride was used instead of 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-3-piperidinol hydrochloride and 1-bromo-2-chloro-5-trifluoromethylbenzene was used instead of 1-bromotoluene in Step 1 thereof, 123 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.87 (m, 3H), 7.45 (m, 3H), 7.17 (m, 2H), 7.00 (m, 2H), 6.20 (s, 1H), 4.69 (m, 1H), 3.58 (m, 2H), 3.17 (m, 2H), 2.30 (m, 2H), 2.25 (s, 3H), 2.01 (t, 2H), 1.73 (m, 4H), 1.47-1.37 (m, 3H)

#### Example 133:

Preparation of

4-[N-[1-[N-(3-methyl-benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that 1-[N-(3-methyl-benzofuran-2-ylcarbonyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 250 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.09 (d, 1H), 7.64 (m, 2H), 7.51-7.32 (m, 6H), 6.73 (s, 1H), 4.70 (m, 1H), 3.78 (d, 1H), 3.63 (d, 1H), 3.41 (m, 1H), 3.29 (td, 1H), 3.17 (s, 3H), 2.78 (m, 1H), 2.61 (s, 3H), 2.27 (m, 2H), 1.95-1.82 (m, 3H), 1.72 (m, 2H), 1.48 (m, 2H), 1.32 (m, 1H)

#### Example 134:

Preparation of

4-[N-[1-[N-(3-methyl-benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(5-fluoro-2-trifluoromethylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 85, except that 4-[N-[1-[N-(3-methyl-benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-3-piperidinol hydrochloride was used instead of 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-3-piperidinol hydrochloride and 1-bromo-2-trifluoromethyl-5-fluorobenzene was used instead of 1-bromotoluene in Step 1 thereof, 128 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.63 (d, 1H), 7.48-7.31 (m, 7H), 6.75 (s, 1H), 4.68 (m, 1H), 3.58 (d, 1H), 3.48 (d, 1H), 3.46 (d, 1H), 3.23 (m, 2H), 2.75 (m, 1H), 2.61 (s, 3H), 2.33(d, 2H), 2.04-1.97 (t, 2H), 1.81-1.72 (m, 4H), 1.52 (m, 2H), 1.21 (m, 1H)

#### Example 135:

Preparation of

4-[N-[1-[N-(3-methyl-benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(5-cyano-2-methoxyphenyl)piperidin-3-one

In accordance with the same procedure as in Example 85, except that 4-[N-[1-[N-(3-methyl-benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-3-piperidinol hydrochloride was used instead of 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-3-piperidinol hydrochloride and 3-bromo-4-methoxybenzonitrile was used instead of 1-bromotoluene in Step 1 thereof, 270 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.63 (d, 1H), 7.50 (m, 3H), 7.33-7.29 (m, 2H), 7.11 (s, 1H), 6.88 (d, 1H), 6.74 (s, 1H), 4.71 (m, 1H), 3.89 (s, 3H), 3.84 (d, 1H), 3.59 (d, 1H), 3.51 (m, 1H), 3.24 (t, 1H), 2.78 (m, H), 2.61 (s, 3H), 2.30 (m, 2H), 2.03 (t, 2H), 1.82-1.61 (m, 4H), 1.20 (m, 2H), 1.18 (m, 1H)

#### Example 136:

Preparation of

4-[N-[1-[N-(quinolin-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that 1-[N-(quinolin-2-ylcarbonyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 310 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.64 (s, 1H), 8.36 (d, 1H), 8.30 (d, 1H), 8.17 (d, 1H), 8.06 (d, 1H), 7.91 (d, 1H), 7.81 (d, 1H), 7.67 (d, 1H), 7.42 (m, 1H), 7.68 (m, 3H), 7.40 (m, 2H), 7.32 (s, 1H), 4.74 (m, 1H), 3.78 (d, 1H), 3.63 (d, 1H), 3.60 (d, 1H), 3.22 (d, 1H), 3.16 (s, 3H), 2.62 (m, 1H), 2.40 (t, 2H), 2.02 (t, 2H), 1.91 (m, 1H), 1.78 (m, 3H), 1.55 (q, 2H), 1.26 (m, 1H)

#### Example 137:

Preparation of

4-[N-[1-[N-(quinolin-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(pyrrolidin-2-one-1-yl)phenyl]piperidin-3-one

In accordance with the same procedure as in Example 92, except that

4-[N-[1-[N-(quinolin-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(4-chlorobutyryl)aminophenyl]piperidin-3-ol was used instead of 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(3-chloro-2,2-dimethylpropionyl)aminophenyl]piperidin-3-ol in Step 2 thereof, 900 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.65 (s, 1H), 8.36 (d, 1H), 8.30 (s, 1H), 8.19 (d, 1H), 7.92 (d, 1H), 7.82 (m, 1H), 7.66 (m, 1H), 7.58 (m, 1H), 7.24 (m, 2H), 7.10 (m, 2H), 4.65 (m, 1H), 3.69 (m, 1H), 3.61 (m, 1H), 3.55 (d, 1H), 3.51 (d, 1H), 3.30 (d, H), 3.21 (t, 1H), 2.84 (m, 2H), 2.51 (m, 2H), 2.33 (m, 2H), 1.74 (m, 4H), 1.28 (m, 2H)

#### Example 138:

Preparation of

4-[N-[1-[N-(quinolin-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-nitrophenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that 1-[N-(quinolin-2-ylcarbonyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, and except that 1-fluoro-2-nitrobenzene was used instead of 1-fluorophenyl methane sulfone in Step 5 thereof, 1.7 g of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.62 (s, 1H), 8.32 (d, 1H), 8.28 (d, 1H), 8.16 (d, 1H), 7.88 (d, 1H), 7.80-7.74 (m, 2H), 7.59 (t, 1H), 7.49 (d, 1H), 7.14-1.09 (m, 2H), 4.70 (m, 1H), 3.73 (s, 2H), 3.37 (m, 2H), 2.86 (m, 1H), 2.31 (m, 2H), 2.02 (m, 2H), 1.91 (m, 1H), 1.75 (m, 2H), 1.54 (f, 2H), 1.40 (m, 1H)

#### Example 139:

Preparation of

4-[N-[1-[N-(pyrazin-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that

1-[N-(pyrazin-2-ylcarbonyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 340 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.8 (m, 1H), 7.3-7.1 (m, 4H), 4.7 (m, 1H),  
5 3.7 (dd, 2H), 3.2 (s, 3H), 1.3-2.1 (m, 10H)

#### Example 140:

Preparation of

4-[N-[1-[N-(pyrazin-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-metha  
10 nesulfonylphenyl)piperidin-3-one dihydrochloride

50 ml of saturated hydrogenchloride-ethyl acetate solution was added to  
4-[N-[1-[N-(pyrazin-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-metha  
nesulfonylphenyl)piperidin-3-one (500 mg) obtained in Example 139. The  
15 reaction mixture was stirred for 2 hours at room temperature and filtered to give  
380 mg of the titled compound as white solid.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 9.87 (broad, 1H), 7.8 (m, 1H), 7.3-7.1 (m,  
4H), 4.7 (m, 1H), 3.7 (dd, 2H), 3.2 (s, 3H), 1.3-2.1 (m, 10H)

#### 20 Example 141:

Preparation of

4-[N-[1-[N-(isonicotinylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-metha  
nesulfonylphenyl)piperidin-3-one

25 In accordance with the same procedure as in Example 76, except that  
1-[N-(isonicotinylcarbonyl)amino]cyclohexanecarboxylic acid was used instead  
of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof,  
430 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.8 (d, 2H), 7.8 (d, 2H), 7.2-7.4 (m, 3H),  
30 6.3 (s, 1H), 4.7 (m, 1H), 3.7 (dd, 2H), 3.2 (s, 3H), 1.3-2.1 (m, 10H)

#### Example 142:

## Preparation of

4-[N-[1-[N-(6-chloronicotinylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

5 In accordance with the same procedure as in Example 76, except that 1-[N-(6-chloronicotinylcarbonyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 122 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.63 (d, 1H), 8.09-7.95 (m, 2H), 7.66 (t, 1H), 7.42 (d, 1H), 7.40 (m, 2H), 6.82 (d, 1H), 6.16 (s, 1H), 4.71 (m, 1H), 3.80 (d, 1H), 3.63 (d, 1H), 3.41 (m, 1H), 3.29 (td, 1H), 3.17 (s, 3H), 2.78 (m, 1H), 2.27 (m, 2H), 1.95-1.82 (m, 3H), 1.72 (m, 2H), 1.48 (m, 2H), 1.32 (m, 1H)

Example 143:

## 15 Preparation of

4-[N-[1-[N-(6-chloronicotinylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-nitrophenyl)piperidin-3-one

20 In accordance with the same procedure as in Example 76, except that 1-[N-(6-chloronicotinylcarbonyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, and except that 1-fluoronitrobenzene was used instead of 1-fluorophenyl methane sulfone in Step 5 thereof, 255 mg of the titled compound was prepared.

25 <sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.62 (s, 1H), 7.99 (m, 1H), 7.53 (m, 2H), 7.21 (m, 2H), 6.77 (d, 1H), 6.27 (s, 1H), 4.71 (m, 1H), 3.98 (m, 3H), 3.69 (s, 2H), 3.34 (m, 2H), 2.78 (m, 1H), 2.23 (d, 2H), 1.97 (m, 4H), 1.70 (m, 4H), 1.45-1.21 (m, 3H)

30 Example 144:

## Preparation of

4-[N-[1-[N-(pyridin-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-metha

nesulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that 1-[N-(pyridin-2-ylcarbonyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 1.0 g of the titled compound was prepared.

$^1\text{H-NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  8.6 (m, 2H), 8.4 (s, 1H), 8.1-8.2 (m, 3H), 7.8 (m, 1H), 7.2-7.6 (m, 3H), 4.7 (m, 1H), 3.6 (dd, 2H), 3.2 (s, 3H), 1.3-2.1 (m, 10H)

Example 145:

Preparation of

4-[N-[1-[N-(pyridin-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one hydrochloride

50 ml of saturated hydrogenchloride-ethyl acetate solution was added to 4-[N-[1-[N-(pyridin-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one (500 mg) obtained in Example 144. The reaction mixture was stirred for 2 hours at room temperature and filtered to give 410 mg of the titled compound as white solid.

$^1\text{H-NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  9.32 (broad, 1H), 8.6 (m, 2H), 8.4 (s, 1H), 8.1-8.2 (m, 3H), 7.8 (m, 1H), 7.2-7.6 (m, 3H), 4.7 (m, 1H), 3.6 (dd, 2H), 3.2 (s, 3H), 1.3-2.1 (m, 10H)

Example 146:

Preparation of

4-[N-[1-[N-(4-fluorobenzoyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that 1-[N-(4-fluorobenzoyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof,



653 mg of the titled compound was prepared.

$^1\text{H-NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  7.8 (m, 2H), 7.6 (m, 1H), 7.4 (m, 2H), 6.1 (s, 1H), 4.7 (m, 1H), 3.7 (dd, 2H), 3.2 (s, 3H), 2.3 (m, 2H), 1.3-2.1 (m, 10H)

5 Example 147:

Preparation of

4-[N-[1-[N-(4-fluorobenzoyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-methanesulfonylphenyl)piperidin-3-one

10 In accordance with the same procedure as in Example 76, except that 1-[N-(4-fluorobenzoyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, and 2,5-difluorophenyl methyl sulfone was used instead of 1-fluorophenyl methane sulfone in Step 5 thereof, 680 mg of the titled compound was  
15 prepared.

$^1\text{H-NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  8.2 (m, 1H), 7.81 (m, 3H), 7.61 (m, 1H), 7.41 (m, 1H), 7.24 (m, 1H), 7.15 (m, 1H), 6.15 (s, 1H), 4.63 (s, 1H), 3.21 (s, 3H), 2.1-1.31(m, 12H)

20 Example 148:

Preparation of

4-[N-[1-[N-(4-fluorobenzoyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonyl-4-methoxyphenyl)piperidin-3-one

25 In accordance with the same procedure as in Example 76, except that 1-[N-(4-fluorobenzoyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, and 2-fluoro-5-methoxyphenyl methyl sulfone was used instead of 1-fluorophenyl methane sulfone in Step 5 thereof, 885 mg of the titled  
30 compound was prepared.

$^1\text{H-NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  8.1 (m, 1H), 7.78 (m, 2H), 7.5 (m, 1H), 7.38 (m, 1H), 7.20 (m, 1H), 7.15 (m, 1H), 6.2 (s, 1H), 4.62 (s, 1H), 3.82 (s, 3H), 3.21

(s, 3H), 2.1-1.31 (m, 12H)

Example 149:

Preparation of

5 4-[N-[1-[N-(4-fluorobenzoyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-trifluoromethylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 85, except that  
10 4-[N-[1-[N-(4-fluorobenzoyl)amino]cyclohexanecarbonyl]amino]-3-piperidinol hydrochloride was used instead of 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-3-piperidinol hydrochloride and 1-bromo-4-fluoro-2-trifluoromethylbenzene was used instead of 1-bromo-4-fluoro-2-thiomethoxybenzene in Step 1 thereof, 137 mg of the titled compound was prepared.

15  $^1\text{H-NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  8.1 (m, 1H), 7.78 (m, 2H), 7.5 (m, 1H), 7.38 (m, 1H), 7.20 (m, 1H), 7.15-6.90 (m, 3H), 6.2 (s, 1H), 4.62 (s, 1H), 2.1-1.31 (m, 12H)

20 Example 150:

Preparation of

4-[N-[1-[N-(4-fluorobenzoyl)amino]cyclohexanecarbonyl]amino]-1-[2-(3-phenylpropionylamino)phenyl]piperidin-3-one

25 In accordance with the same procedure as in Example 87, except that 4-[N-[1-[N-(4-fluorobenzoyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-aminophenyl)piperidin-3-ol was used instead of 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-aminophenyl)piperidin-3-ol and hydrocinnamoyl chloride was used instead of methoxyacetyl chloride in Step 3 thereof, 671 mg of the titled compound was  
30 prepared.

$^1\text{H-NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  8.1 (m, 1H), 7.78 (m, 2H), 7.5 (m, 1H),

7.38 (m, 1H), 7.20 (m, 1H), 7.15-6.90 (m, 8H), 6.2 (s, 1H), 4.62 (s, 1H), 2.83 (m, 2H), 2.44 (m, 2H), 2.19 (m, 2H), 2.1-1.31 (m, 12H)

5 Example 151:

Preparation of

4-[N-[1-[N-(4-fluorobenzoyl)amino]cyclohexanecarbonyl]amino]-1-[2-(cyclopropylamino)phenyl]piperidin-3-one

10 In accordance with the same procedure as in Example 87, except that 4-[N-[1-[N-(4-fluorobenzoyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-aminophenyl)piperidin-3-ol was used instead of 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-aminophenyl)piperidin-3-ol and cyclopropylcarbonyl chloride was used instead  
15 of methoxyacetyl chloride in Step 3 thereof, 1.5 g of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.1 (m, 1H), 7.78 (m, 2H), 7.5 (m, 1H), 7.38 (m, 1H), 7.20 (m, 1H), 7.15-6.90 (m, 3H), 6.2 (s, 1H), 4.62 (s, 1H), 2.1-1.31 (m, 12H), 0.48 (m, 4H)

20

Example 152:

Preparation of

4-[N-[1-[N-(4-fluorobenzoyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(2,3,4-trifluorobenzoylamino)phenyl]piperidin-3-one  
25

In accordance with the same procedure as in Example 87, except that 4-[N-[1-[N-(4-fluorobenzoyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-aminophenyl)piperidin-3-ol was used instead of  
30 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-aminophenyl)piperidin-3-ol and 2,3,4-trifluorobenzoyl chloride was used instead of methoxyacetyl chloride in Step 3 thereof, 321 mg of the titled compound was

prepared.

$^1\text{H-NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  8.1 (m, 1H), 7.78 (m, 2H), 7.5 (m, 1H), 7.38 (m, 1H), 7.20 (m, 1H), 7.15-6.90 (m, 3H), 6.2 (s, 1H), 4.62 (s, 1H), 2.1-1.31 (m, 12H)

5

Example 153:

Preparation of

4-[N-[1-[N-(4-biphenylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

10

In accordance with the same procedure as in Example 76, except that 1-[N-(4-biphenylcarbonyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 653 mg of the titled compound was prepared.

15

$^1\text{H-NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  8.11 (d, 1H), 7.90 (d, 2H), 7.66 (m, 5H), 7.38 (m, 5H), 4.71 (m, 1H), 3.78 (d, 1H), 3.75 (d, 1H), 3.64 (d, 1H), 3.56(d, 1H), 3.19 (s, 3H), 2.78 (s, 3H), 2.15 (m, 2H), 1.89 (m, 3H), 1.66 (m, 4H), 1.38-1.26 (m, 3H)

20

Example 154:

Preparation of

4-[N-[1-[N-(4-trifluoromethylbenzoyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

25

In accordance with the same procedure as in Example 76, except that 1-[N-(4-trifluoromethylbenzoyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 432 mg of the titled compound was prepared.

30

$^1\text{H-NMR}$  (400MHz,  $\text{CDCl}_3$ )  $\delta$  8.10 (d, 1H), 8.07 (d, 2H), 7.87 (d, 1H), 7.68 (d, 1H), 7.48 (d, 1H), 7.38 (m, 2H), 7.29 (m, 2H), 6.20 (s, 1H), 4.71 (m, 1H), 3.78 (d, 1H), 3.64 (d, 1H), 3.20 (s, 3H), 2.78 (m, 1H), 2.05 (m, 2H),

1.87 (m, 3H), 1.62 (m, 5H), 1.16 (m, 3H)

Example 155:

5 Preparation of

4-[N-[1-[N-(4-(2-morpholinethoxy)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

10 In accordance with the same procedure as in Example 76, except that 1-[N-(4-(2-morpholinethoxy)amino)cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 543 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.05 (d, 1H), 7.8 (d, 2H), 7.68-7.24 (m, 4H), 6.68 (d, 2H), 4.89 (m, 1H), 4.08 (m, 4H), 3.75 (t, 2H), 3.15 (s, 3H), 2.18-1.15 (m, 15 12H)

Example 156:

Preparation of

20 4-[N-[1-[N-(4-(2-pyrrolidinethoxy)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

25 In accordance with the same procedure as in Example 76, except that 1-[N-(4-(2-pyrrolidinethoxy)benzoyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 543 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.05 (d, 1H), 7.8 (d, 2H), 7.68-7.24 (m, 4H), 6.68 (d, 2H), 4.69 (m, 1H), 4.08 (m, 4H), 3.82 (dd, 2H), 3.15 (s, 3H), 2.18-1.15 (m, 12H)

30

Example 157:

Preparation of

4-[N-[1-[N-(4-(2-piperidinethoxy)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that  
5 1-[N-(4-(2-piperidinethoxy)benzoyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 543 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.05 (d, 1H), 7.8 (d, 2H), 7.68-7.24 (m, 4H),  
6.68 (d, 2H), 4.69 (m, 1H), 4.08 (m, 4H), 3.82 (dd, 2H), 3.15 (s, 3H), 2.8 (m, 2H),  
10 2.18-1.15 (m, 12H)

Example 158:

Preparation of

15 4-[N-[1-[N-(4-(2-morpholinethoxy)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one hydrochloride

50 ml of saturated hydrogenchloride-ethyl acetate solution was added to  
4-[N-[1-[N-(4-(2-morpholinethoxy)benzoyl)amino]cyclohexanecarbonyl]amino]-1  
20 -(2-methanesulfonylphenyl)piperidin-3-one (500 mg) obtained in Example 155. The reaction mixture was stirred for 2 hours at room temperature and filtered to give 420 mg of the titled compound as white solid.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.05 (d, 1H), 7.83 (m, 5H), 7.52 (m, 2H),  
7.24 (m, 2H), 6.88 (m, 2H), 4.65 (m, 4H), 4.2 (m, 4H), 3.92 (m, 2H), 3.62-3.21  
25 (m, 5H), 2.1-1.31 (m, 12H)

Example 159:

Preparation of

30 4-[N-[1-[N-(4-(2-pyrrolidinethoxy)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one hydrochloride

50 ml of saturated hydrogenchloride-ethyl acetate solution was added to 4-[N-[1-[N-(4-(2-pyrrolidinethoxy)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one (500 mg) obtained in Example 156. The reaction mixture was stirred for 2 hours at room temperature and filtered to give 320 ml of the titled compound as white solid.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.05 (d, 1H), 7.8-7.52 (m, 7H), 7.24 (m, 2H), 6.68 (m, 2H), 4.89 (m, 1H), 2.85-3.6 (m, 8H), 2.18-1.15 (m, 12H)

Example 160:

10 Preparation of  
4-[N-[1-[N-(4-(2-piperidinethoxy)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one hydrochloride

50 ml of saturated hydrogenchloride-ethyl acetate solution was added to 4-[N-[1-[N-(4-(2-piperidinethoxy)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one (500 mg) obtained in Example 157. The reaction mixture was stirred for 2 hours at room temperature and filtered to give 472 mg of the titled compound as white solid.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.05 (d, 1H), 7.8-7.52 (m, 7H), 7.24 (m, 2H), 6.68 (m, 2H), 4.89 (m, 1H), 2.85-3.6 (m, 8H), 2.25 (m, 2H), 2.18-1.15 (m, 12H)

Example 161:

25 Preparation of  
4-[N-[1-[N-(5-(2-morpholinethoxy)benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that 1-[N-(5-(2-morpholinethoxy)benzofuran-2-ylcarbonyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof,

543 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.05 (m, 1H), 7.8 (m, 2H), 7.68-7.24 (m, 4H), 6.68 (m, 2H), 4.69 (m, 1H), 4.08 (m, 4H), 3.82-3.4 (m, 4H), 3.15 (s, 3H), 2.8 (m, 2H), 2.18-1.15 (m, 12H)

5

Example 162:

Preparation of

4-[N-[1-[N-(5-(2-pyrrolidinethoxy)benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

10

In accordance with the same procedure as in Example 76, except that 1-[N-(5-(2-pyrrolidinethoxy)benzofuran-2-ylcarbonyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 759 mg of the titled compound was prepared.

15

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.05 (m, 1H), 7.8 (m, 2H), 7.68-7.24 (m, 4H), 6.68 (m, 2H), 4.69 (m, 1H), 4.2 (m, 2H), 3.82-3.6 (dd, 2H), 3.15 (s, 3H), 2.8 (m, 2H), 2.18-1.15 (m, 12H)

20

Example 163:

Preparation of

4-[N-[1-[N-(5-(2-piperidinethoxy)benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

25

In accordance with the same procedure as in Example 76, except that 1-[N-(5-(2-piperidinethoxy)benzofuran-2-ylcarbonyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 990 mg of the titled compound was prepared.

30

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.05 (m, 1H), 7.8 (m, 2H), 7.68-7.24 (m, 4H), 6.68 (m, 2H), 4.69 (m, 1H), 4.2 (m, 2H), 3.82-3.6 (dd, 2H), 3.15 (s, 3H), 2.8 (m, 2H), 2.18-1.15 (m, 14H)



Example 164:

## Preparation of

5 4-[N-[1-[N-(5-(2-morpholinethoxy)benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one hydrochloride

50 ml of saturated hydrogenchloride-ethyl acetate solution was added to 4-[N-[1-[N-(5-(2-morpholinethoxy)benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one (500 mg) obtained in Example 161. The reaction mixture was stirred for 2 hours at room temperature and filtered to give 472 mg of the titled compound as white solid.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.05 (d, 1H), 7.8 (d, 2H), 7.68-7.24 (m, 4H), 6.68 (d, 2H), 4.69 (m, 1H), 4.08 (m, 4H), 3.88 (m, 2H), 3.65 (m, 4H), 3.15-3.55 (m, 7H), 2.8 (m, 2H), 2.18-1.15 (m, 12H)

Example 165:

## Preparation of

20 4-[N-[1-[N-(5-(2-pyrrolidinethoxy)benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one hydrochloride

50 ml of saturated hydrogenchloride-ethyl acetate solution was added to 4-[N-[1-[N-(5-(2-pyrrolidinethoxy)benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one (500 mg) obtained in Example 162. The reaction mixture was stirred for 2 hours at room temperature and filtered to give 130 mg of the titled compound as white solid.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.05 (d, 1H), 7.8 (d, 2H), 7.68-7.24 (m, 4H), 6.68 (d, 2H), 4.69 (m, 1H), 4.08 (m, 4H), 3.82 (dd, 2H), 3.15 (s, 3H), 2.18-1.15 (m, 12H)

Example 166:

## Preparation of

4-[N-[1-[N-(5-(2-piperidinethoxy)benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one hydrochloride

5            50 ml of saturated hydrogenchloride-ethyl acetate solution was added to 4-[N-[1-[N-(5-(2-piperidinethoxy)benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one (500 mg) obtained in Example 163. The reaction mixture was stirred for 2 hours at room temperature and filtered to give 250 mg of the titled compound as white solid.

10            <sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.05 (d, 1H), 7.68 (m, 1H), 7.66-7.34 (m, 4H), 7.11 (s, 1H), 6.68 (d, 2H), 4.69 (m, 1H), 4.48 (m, 1H), 3.82-3.34 (m, 8H), 3.15 (s, 3H), 2.8 (m, 2H), 2.18-1.15 (m, 12H)

15            Example 167:

## Preparation of

4-[N-[1-[N-(4-(2-oxopyrrolidine)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

20            In accordance with the same procedure as in Example 76, except that 1-[N-(4-(2-oxopyrrolidine)benzoyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 770 mg of the titled compound was prepared.

25            <sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.05 (m, 1H), 7.78-7.24 (m, 8H), 6.65 (m, 1H), 4.69 (m, 1H), 3.92 (m, 2H), 3.82-3.6 (dd, 2H), 3.15 (s, 3H), 2.8 (m, 2H), 2.2-2.42 (m, 4H), 2.18-1.15 (m, 12H)

Example 168:

## 30            Preparation of

4-[N-[1-[N-(4-(2-oxopyrrolidine)furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that 1-[N-(4-(2-oxopyrrolidine)furan-2-ylcarbonyl)amino]cyclohexane carboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 770 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.05 (m, 1H), 7.65 (m, 1H), 7.38-7.14 (m, 3H), 6.48 (m, 1H), 4.69 (m, 1H), 4.2 (m, 2H), 3.82-3.6 (dd, 2H), 3.15 (s, 3H), 2.8 (m, 2H), 2.18~1.15(m,12H)

#### Example 169:

Preparation of

4-[N-[1-[N-(4-(2-morpholinethoxy)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-methanesulfonylaminophenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that 1-[N-(4-(2-morpholinethoxy)benzoyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof and 2,5-difluoro-1-methanesulfonylbenzene was used instead of 1-fluorophenyl methyl sulfone in Step 5 thereof, 155 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.05 (d, 1H), 7.83 (m, 5H), 7.52 (m, 2H), 7.24 (m, 2H), 6.88 (m, 2H), 4.65 (m, 4H), 4.2 (m, 4H), 3.92 (m, 2H), 3.62-3.21 (m, 5H), 2.1-1.31 (m, 12H)

#### Example 170:

Preparation of

4-[N-[1-[N-(4-(2-morpholinethoxy)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-(pyrrolidin-2-one-1-yl)phenyl)piperidin-3-one

In accordance with the same procedure as in Example 92, except that

4-[N-[1-[N-(4-(2-morpholinethoxy)benzoyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(4-chlorobutyryl)aminophenyl]piperidin-3-ol was used instead of 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(3-chloro-2,2-dimethylpropionyl)aminophenyl]piperidin-3-ol in Step 2 thereof, 130 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.05 (d, 1H), 7.83 (m, 5H), 7.52 (m, 2H), 7.24 (m, 2H), 6.88 (m, 2H), 4.65 (m, 4H), 4.2 (m, 4H), 3.92 (m, 2H), 3.62-3.21 (m, 5H), 2.1-1.31 (m, 16H)

#### Example 171:

Preparation of

4-[N-[1-[N-(4-(dimethylamino)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-methanesulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that 1-[N-(4-(dimethylamino)benzoyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof and 2,5-difluorophenyl methane sulfone was used instead of 1-fluorophenyl methyl sulfone in Step 5 thereof, 222 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.05 (d, 1H), 7.83 (m, 5H), 7.52 (m, 2H), 7.24 (m, 2H), 6.68 (d, 2H), 6.01 (s, 1H), 4.65 (m, 1H), 4.65-4.79 (dd, 2H), 3.25 (s, 3H), 3.05 (s, 6H), 2.1-1.31 (m, 12H)

#### Example 172:

Preparation of

4-[N-[1-[N-(4-(diethylamino)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-methanesulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that

1-[N-(4-(diethylamino)benzoyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof and 2,5-difluorophenyl methane sulfone was used instead of 1-fluorophenyl methane sulfone in Step 5 thereof, 332 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.05 (d, 1H), 7.83 (m, 5H), 7.52 (m, 2H), 7.24 (m, 2H), 6.68 (d, 2H), 6.01 (s, 1H), 4.68 (m, 1H), 4.65-4.79 (dd, 2H), 3.35 (m, 4H), 3.21 (s, 3H), 2.2-1.15 (m, 12H), 1.05 (m, 6H)

#### Example 173:

Preparation of

4-[N-[1-[N-(4-(dimethylamino)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-methanesulfonylphenyl)piperidin-3-one hydrochloride

50 ml of saturated hydrogenchloride-ethyl acetate solution was added to 4-[N-[1-[N-(4-(dimethylamino)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-methanesulfonylphenyl)piperidin-3-one (500 mg) obtained in Example 171. The reaction mixture was stirred for 2 hours at room temperature and filtered to give 250 mg of the titled compound as white solid.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.0-8.2 (m, 2H), 7.8-7.6 (m, 2H), 7.48 (m, 1H), 7.38 (m, 2H), 4.68 (m, 1H), 3.28 (s, 3H), 3.02-3.2 (m, 6H), 3.21 (s, 3H), 2.18-1.31 (m, 12H)

#### Example 174:

Preparation of

4-[N-[1-[N-(4-(diethylamino)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-methanesulfonylphenyl)piperidin-3-one hydrochloride

50 ml of saturated hydrogenchloride-ethyl acetate solution was added to 4-[N-[1-[N-(4-(diethylamino)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-methanesulfonylphenyl)piperidin-3-one (500 mg) obtained in Example

172. The reaction mixture was stirred for 2 hours at room temperature and filtered to give 440 mg of the titled compound as white solid.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.0-8.2(m, 2H), 7.8(m, 1H), 7.52(m, 2H), 7.41(m, 2H), 7.15 (s, 1H), 4.72 (s, 1H), 3.4 (m, 4H), 3.21 (s, 3H), 2.18-1.31(m, 12H), 1.1(m, 6H)

#### Example 175:

Preparation of

4-[N-[1-[N-(4-(3-phenylpropionyl)amino)cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that 1-[N-(4-(3-phenylpropionyl)amino)cyclohexane carboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 132 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.63 (d, 1H), 7.48-7.00 (m, 8H), 5.10 (s, 2H), 4.68 (m, 1H), 3.58 (d, 1H), 3.48 (d, 1H), 3.46 (d, 1H), 3.23 (m, 2H), 2.99 (s, 3H), 2.75 (m, 3H), 2.33 (m, 4H), 2.04-1.97 (t, 2H), 1.81-1.72 (m, 4H), 1.52 (m, 2H), 1.21 (m, 1H)

#### Example 176:

Preparation of

4-[N-[1-[N-(2-methylcinnamoyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that 1-[N-(2-methylcinnamoyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 450 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.10 (d, 1H), 7.68 (d, 1H), 7.34 (m, 8H), 5.99 (s, 1H), 4.71 (m, 1H), 4.09 (m, 1H), 3.78 (d, 1H), 3.67 (d, 1H), 3.63 (d, 1H), 3.20 (s, 3H), 2.78 (m, 1H), 2.25 (m, 2H), 2.15 (s, 3H), 1.89 (m, 3H),

1.66 (m, 4H), 1.38-1.26 (m, 3H)

Example 177:

Preparation of

5 4-[N-[1-[N-(benzyloxycarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-acetylamino)phenyl]piperidin-3-one

In accordance with the same procedure as in Example 87, except that  
4-[N-[1-[N-(benzyloxycarbonyl)amino]cyclohexanecarbonyl]amino]-3-piperidinol  
10 hydrochloride was used instead of  
4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-3-piperidinol  
hydrochloride in Step 1 thereof, and acetyl chloride was used instead of  
methoxyacetyl chloride in Step 3 of thereof, 670 mg of the titled compound was  
prepared.

15 <sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.56 (d, 1H), 7.48-6.95 (m, 8H), 5.10 (s, 2H), 4.68 (m, 1H), 3.58 (d, 1H), 3.48 (d, 1H), 3.46 (d, 1H), 3.23 (m, 2H), 2.75 (m, 1H), 2.33 (d, 2H), 2.04-1.97 (t, 2H), 1.81-1.72 (m, 4H), 1.52 (m, 2H), 1.21 (m, 1H)

20 Example 178:

Preparation of

4-[N-[1-[N-(benzyloxycarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-trifluoromethylphenyl)piperidin-3-one

25 In accordance with the same procedure as in Example 24, except that  
4-[N-[1-[N-(benzyloxycarbonyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-aminophenyl)piperidin-3-ol hydrochloride was used instead of  
4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-3-piperidinol hydrochloride and 1-bromo-2-trifluoromethylbenzene was used instead of  
30 1-bromo-4-fluoro-2-trifluoromethylbenzene in Step 1 of Example 24, 228 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 7.63 (d, 1H), 7.48-7.00 (m, 8H), 5.10 (s,

2H), 4.68 (m, 1H), 3.58 (d, 1H), 3.48 (d, 1H), 3.46 (d, 1H), 3.23 (m, 2H), 2.75 (m, 1H), 2.33 (d, 2H), 2.04-1.97 (t, 2H), 1.81-1.72 (m, 4H), 1.52 (m, 2H), 1.21 (m, 1H)

5 Example 179:

Preparation of

4-[N-[1-[N-(pyran-2-one-5-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

10 In accordance with the same procedure as in Example 76, except that 1-[N-(pyran-2-one-5-ylcarbonyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 750 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.62 (broad, 1H), 8.10 (d, 1H), 7.67 (m, 2H), 7.38 (m, 2H), 7.30 (s, 1H), 7.20 (d, 1H), 6.89 (d, 1H), 5.71 (d, 1H), 4.71 (m, 1H), 4.09 (m, 1H), 3.78 (d, 1H), 3.67 (d, 1H), 3.63 (d, 1H), 2.78 (m, 1H), 2.15 (m, 2H), 1.89 (m, 3H), 1.66 (m, 4H), 1.38-1.26 (m, 3H),

Example 180:

20 Preparation of

4-[N-[1-[N-((6-methyl)-7H-imidazo[2,1-b]oxazol-5-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

25 In accordance with the same procedure as in Example 76, except that 1-[N-((6-methyl)-7H-imidazo[2,1-b]oxazol-5-ylcarbonyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 59 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.24 (d, 1H), 8.10 (d, 1H), 7.68 (d, 1H), 7.42 (d, 1H), 7.38 (m, 2H), 6.89 (d, 1H), 4.76 (m, 1H), 3.78 (d, 1H), 3.64 (d, 1H), 3.56 (d, 1H), 3.19 (s, 3H), 2.78 (s, 3H), 2.15 (m, 2H), 1.89 (m, 3H), 1.66 (m, 4H), 1.38-1.26 (m, 3H)



Example 181:

Preparation of

4-[N-[1-[N-((6-methyl)-chromon-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]  
5 ]-1-(2-methanesulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that  
1-[N-((6-methyl)-chromon-2-ylcarbonyl)amino]cyclohexanecarboxylic acid was  
used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in  
10 Step 3 thereof, 82 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.54 (broad, 1H), 8.10 (d, 1H), 7.87 (s,  
1H), 7.70 (s, 1H), 7.68 (d, 1H), 7.42 (d, 1H), 7.38 (m, 4H), 7.03 (s, 1H), 6.83 (s,  
1H), 5.19 (m, 1H), 4.32 (m, 1H), 3.78 (d, 1H), 3.64 (d, 1H), 3.56(d, 1H),  
3.19 (s, 3H), 2.78 (s, 3H), 2.15 (m, 2H), 1.89 (m, 3H), 1.66 (m, 4H),  
15 1.38-1.26 (m, 3H)

Example 182:

Preparation of

4-[N-[1-[N-(isobutyryl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonyl  
20 phenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that  
1-[N-(isobutyryl)amino]cyclohexanecarboxylic acid was used instead of  
1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof,  
25 382 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.10 (d, 1H), 7.68 (d, 1H), 7.38 (m, 2H),  
5.52 (s, 1H), 4.66 (m, 1H), 3.78 (d, 1H), 3.64 (d, 1H), 3.36(d, 1H), 3.27 (d, 1H),  
3.19 (s, 3H), 2.78 (m, 1H), 2.44 (m, 1H), 2.17 (t, 2H), 1.89-1.71 (m, 3H),  
1.69-1.49 (m, 3H), 1.34 (m, 3H), 1.18 (m, 7H)

30

Example 183:

Preparation of

4-[N-[1-[N-(isovaleryl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that  
5 1-[N-(isovaleryl)amino]cyclohexanecarboxylic acid was used instead of  
1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof,  
442 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.10 (d, 1H), 7.68 (d, 1H), 7.38 (m, 2H),  
5.52 (s, 1H), 4.66 (m, 1H), 3.78 (d, 1H), 3.64 (d, 1H), 3.36(d, 1H), 3.27 (d, 1H),  
10 3.19 (s, 3H), 2.78 (m, 1H), 2.17 (m, 4H), 1.89-1.71 (m, 3H), 1.69-1.49 (m, 4H),  
1.37 (m, 3H), 0.99 (d, 6H)

Example 184:

Preparation of

15 4-[N-[1-[N-(3,3-dimethylacryloyl)amino]cyclohexanecarbonyl]amino]-1-(2-metha  
nesulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that  
20 1-[N-(3,3-dimethylacryloyl)amino]cyclohexanecarboxylic acid was used instead  
of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof,  
442 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.10 (d, 1H), 7.68 (d, 1H), 7.38 (m, 2H),  
5.06 (d, 1H, J = 28Hz), 4.66 (m, 1H), 3.78 (d, 1H), 3.64 (d, 1H), 3.52 (d, 1H),  
3.20 (s, 3H), 2.78 (m, 1H), 2.17 (m, 2H), 1.89 (m, 6H), 1.69-1.49 (m, 5H), 1.37  
25 (m, 3H),

Example 185:

Preparation of

4-[N-[1-[N-(2-methoxyacetyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanes  
30 ulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that

1-[N-(2-methoxyacetyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 872 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.10 (d, 1H), 7.68 (d, 1H), 7.38 (m, 2H),  
5 6.68 (s, 1H), 4.66 (m, 1H), 3.94 (s, 2H), 3.78 (d, 1H), 3.64 (d, 1H), 3.49 (s, 3H),  
3.27 (d, 1H), 3.19 (s, 3H), 2.78 (m, 1H), 2.17 (m, 2H), 1.89-1.71 (m, 4H),  
1.69-1.49 (m, 4H), 1.37 (m, 4H)

Example 186:

10 Preparation of

4-[N-[1-[N-(cyclopropanecarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that  
15 1-[N-(cyclopropanecarbonyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 731 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.10 (d, 1H), 7.68 (d, 1H), 7.38 (m, 2H),  
5.74 (s, 1H), 4.71 (m, 1H), 3.78 (d, 1H), 3.64 (d, 1H), 3.20 (s, 3H), 2.78 (m,  
20 1H), 2.12 (m, 2H), 1.89 (m, 3H), 1.66 (m, 4H), 1.43 (m, 4H), 0.99 (m, 2H),  
0.78 (m, 2H)

Example 187:

Preparation of

25 4-[N-[1-[N-(cyclobutanecarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that  
1-[N-(cyclobutanecarbonyl)amino]cyclohexanecarboxylic acid was used instead  
30 of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 639 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.10 (d, 1H), 7.68 (d, 1H), 7.38 (m, 2H),

5.52 (s, 1H), 4.66 (m, 1H), 3.78 (d, 1H), 3.64 (d, 1H), 3.36(d, 1H), 3.19 (s, 3H), 3.08 (m, 1H), 2.78 (m, 1H), 2.32 (m, 2H), 2.17 (m, 4H), 1.89-1.71 (m, 5H), 1.69-1.49 (m, 5H), 1.37 (m, 3H)

5 Example 188:

Preparation of

4-[N-[1-[N-(cyclopentanecarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

10 In accordance with the same procedure as in Example 76, except that 1-[N-(cyclopentanecarbonyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 888 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.10 (d, 1H), 7.68 (d, 1H), 7.38 (m, 2H),  
15 5.57 (s, 1H), 4.66 (m, 1H), 3.78 (d, 1H), 3.64 (d, 1H), 3.36(d, 1H), 3.19 (s, 3H), 3.08 (m, 1H), 2.78 (m, 1H), 2.22-2.15 (m, 2H), 1.92-1.58 (m, 16H), 1.35 (m, 4H)

Example 189:

20 Preparation of

4-[N-[1-[N-(cyclohexanecarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that  
25 1-[N-(cyclohexanecarbonyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 565 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.10 (d, 1H), 7.68 (d, 1H), 7.38 (m, 2H),  
5.57 (s, 1H), 4.66 (m, 1H), 3.78 (d, 1H), 3.64 (d, 1H), 3.36(d, 1H), 3.19 (s, 3H),  
30 3.08 (m, 1H), 2.78 (m, 1H), 2.22-2.15 (m, 3H), 1.92-1.71 (m, 8H), 1.69-1.49 (m, 3H), 1.37-1.24 (m, 11H)

Example 190:

Preparation of

4-[N-[1-[N-((3-isobutyloxy)propionyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

5

In accordance with the same procedure as in Example 76, except that 1-[N-((3-isobutyloxy)propionyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 777 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.10 (d, 1H), 7.68 (d, 1H), 7.38 (m, 2H), 6.76 (s, 1H), 4.66 (m, 1H), 3.78 (m, 3H), 3.71 (d, 1H), 3.64(d, 1H), 3.31 (t, 2H), 3.20 (s, 3H), 3.08 (m, 1H), 2.78 (m, 1H), 2.57 (t, 2H), 2.22-2.15 (m, 2H), 1.92-1.72 (m, 4H), 1.66-1.51 (m, 4H), 1.40 (m, 3H), 0.99 (d, 6H)

15 Example 191:

Preparation of

4-[N-[1-[N-((3-benzyloxy)propionyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

20 In accordance with the same procedure as in Example 76, except that 1-[N-((3-benzyloxy)propionyl)amino]cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 337 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.62 (s, 1H), 8.10 (d, 1H), 7.6 (m, 2H), 7.38 (m, 7H), 6.60 (s, 1H), 4.63 (m, 3H), 3.82 (t, 2H), 3.78 (m, 3H), 3.71 (d, 1H), 3.64(d, 1H), 3.31 (t, 2H), 3.20 (s, 3H), 2.70 (m, 1H), 2.57 (t, 2H), 2.22-2.15 (m, 2H), 1.92-1.72 (m, 4H), 1.66-1.51 (m, 4H), 1.40 (m, 3H), 1.22 (d, 3H)

Example 192:

30 Preparation of

4-[N-[1-[N-((N-t-butoxycarbonyl)piperidin-4-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one

In accordance with the same procedure as in Example 76, except that 1-[N-((N-t-butoxyacarbonyl)piperidin-4-ylcarbonyl)amino] cyclohexanecarboxylic acid was used instead of 1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarboxylic acid in Step 3 thereof, 442 mg of the titled compound was prepared.

<sup>1</sup>H-NMR (400MHz, CDCl<sub>3</sub>) δ 8.10 (d, 1H), 7.68 (d, 1H), 7.38 (m, 2H), 5.55 (s, 1H), 4.96 (m, 1H), 4.19 (m, 2H), 3.97 (m, 3H), 3.38 (d, 1H), 3.29 (s, 3H), 3.18 (d, 1H), 2.81 (m, 6H), 2.61 (t, 1H), 2.56 (t, 1H), 2.17 (m, 6H), 1.82 (m, 6H), 1.57 (m, 6H), 1.45 (s, 9H), 1.30 (m, 3H) 2.78 (m, 1H), 2.22-2.15 (m, 2H), 1.92-1.58 (m, 16H), 1.35 (m, 4H)

#### Test 1: Inhibition of cathepsin K activity

Recombinant human cathepsin K was prepared in accordance with *Biol. Pharm. Bull.* 19(8), 1026-1031(1996) and *J. Biol. Chem.* 271, 2126-2132(1996).

Stock solutions of 2.5 mM Z-Phe-Arg-AMC (BACHEM, I-1160) and all inhibitors were prepared in dimethyl sulfoxide. *In vitro* assays for cathepsin K were conducted with 25 uM Z-Phe-Arg-AMC as a substrate at pH 5.5 in 100 mM Na acetate buffer containing 20 mM cysteine and 5 mM EDTA in the presence of 2 % final concentration of DMSO. Reaction was initiated by the addition of enzyme to samples, which were then incubated for 1 hour at 37 °C. The fluorescence was monitored at 460 nm (excitation wavelength 360 nm) using fluorescence plate reader.

The concentrations of the test compounds, which inhibit 50% of cathepsin K activity, are represented as IC<sub>50</sub> in Table 1.

Table 1

	Test Compound	IC <sub>50</sub> (nM)	Test Compound	IC <sub>50</sub> (nM)
5	Example 1	8.17	Example 80	5.05
	Example 2	5.97	Example 83	2.69
	Example 5	5.05	Example 84	6.24
	Example 6	2.69	Example 85	10.4
	Example 7	6.24	Example 92	1.45
10	Example 8	10.4	Example 95	10.8
	Example 15	1.45	Example 99	4.41
	Example 16	10.8	Example 106	19.9
	Example 19	4.41	Example 107	8.07
15	Example 26	19.9	Example 114	1.93
	Example 27	8.07	Example 116	2.35
	Example 34	1.93	Example 117	2.13
	Example 35	2.35	Example 121	7.73
20	Example 36	2.13	Example 122	3.87
	Example 37	3.78	Example 123	4.97
	Example 41	7.73	Example 124	4.96
	Example 42	3.87	Example 125	6.80
25	Example 43	4.97	Example 128	0.67
	Example 44	4.96	Example 129	0.45
	Example 45	6.80	Example 130	1.20
	Example 48	0.67	Example 131	4.11
30	Example 49	0.45	Example 133	1.22
	Example 50	1.20	Example 134	6.09
	Example 51	4.11	Example 136	1.62
	Example 53	1.22	Example 138	4.82
35	Example 54	6.09	Example 147	4.77
	Example 56	1.62	Example 148	3.12
	Example 58	4.82	Example 153	0.36
	Example 67	4.77	Example 154	1.37
	Example 68	3.12	Example 155	1.29
40	Example 69	1.59	Example 156	1.90
	Example 70	1.73	Example 158	3.15
	Example 75	0.51	Example 159	3.25
	Example 76	8.17	Example 160	3.14
45	Example 77	5.97	Example 161	0.41

Table 1 (Continued)

Test Compound	IC <sub>50</sub> (nM)	Test Compound	IC <sub>50</sub> (nM)
Example 162	0.45	Example 169	0.58
Example 163	0.55	Example 170	1.69
Example 164	0.52	Example 171	0.77
Example 165	1.63	Example 172	0.58
Example 166	0.71	Example 173	1.59
Example 167	3.07	Example 174	1.00
Example 168	15.79		

As shown in Table 1, the compounds of the present invention have a cathepsin K activity inhibition effect.

#### Test 2: Selectivity Study

Selectivity study for the compounds of the present invention was performed using cathepsin C (Sigma, C8511), cathepsin G (Calbiochem, 219373), cathepsin H (Calbiochem, 219388), cathepsin L (Calbiochem, 219402), and cathepsin S (Eur. J. Biochem. 236, 558 (1996) as enzyme sources.

##### 1) Cathepsin C

Cathepsin C (Sigma, C8511) was dissolved in D.W. buffer (pH 5.5). All inhibitors were prepared in dimethyl sulfoxide (DMSO). *In vitro* assays for cathepsin C were conducted with Gly-Phe-pNA (Sigma, G-0142) as a substrate.

Samples were pre-incubated for 10 minutes at 37 °C. Reaction was initiated by the addition of enzyme to samples, which were then incubated for 1 hour at 37 °C. The absorbance of *p*-nitroaniline released was monitored at 405 nm using spectrophotometer.

##### 2) Cathepsin G



Cathepsin G (Calbiochem, 219373) was dissolved in 50 mM NaOAc and 150 mM NaCl buffer (pH 5.5). All inhibitors were prepared in dimethyl sulfoxide (DMSO). *In vitro* assays for cathepsin G were conducted with  
5 Suc-Ala-Ala-Pro-Phe-pNA (Calbiochem, 219407) as a substrate. Samples were pre-incubated for 10 minutes at 25 °C. Reaction was initiated by the addition of enzyme to samples, which were then incubated for 1 hour at 25 °C. The absorbance of *p*-nitroaniline released was monitored at 405 nm using spectrophotometer.

10

### 3) Cathepsin H

Cathepsin H (Calbiochem, 219388) was dissolved in 20 mM NaOAc, 1 mM EDTA-2Na and 100 mM NaCl buffer (pH 4.6). All inhibitors were prepared in  
15 dimethyl sulfoxide (DMSO). *In vitro* assays for cathepsin H were conducted with *L*-Arg-pNA (BACHEM, I-1160) as a substrate. Samples were pre-incubated for 5 minutes at 40 °C. Reaction was initiated by the addition of enzyme to samples, which were then incubated for 1 hour at 40 °C. The absorbance of *p*-nitroaniline released was monitored at 405 nm using  
20 spectrophotometer.

### 4) Cathepsin L

Cathepsin L (Calbiochem, 219402) was dissolved in 50 mM NaOAc  
25 buffer (pH 5.0). All inhibitors were prepared in dimethyl sulfoxide (DMSO). *In vitro* assays for cathepsin G were conducted with Z-Phe-Arg-AMC (BACHEM, I-1160) as a substrate. Samples were pre-incubated for 5 minutes at 25 °C. Reaction was initiated by the addition of enzyme to samples, which were then incubated for 1 hour at 25 °C. The fluorescence of AMC

(7-amino-4-methylcoumarin) was monitored at 460 nm (excitation wavelength 360nm), using fluorescence spectrophotometer.

#### 5) Cathepsin S

5

Recombinant human Cathepsin S (Eur. J. Biochem. 236, 558 (1996) was dissolved in 0.1 M NaOAc 1 mM EDTA-4Na buffer (pH 5.5). All inhibitors were prepared in dimethyl sulfoxide (DMSO). *In vitro* assays for cathepsin S were conducted with Z-Val-Val-AMC (BACHEM, I-1160) as a substrate.

10

Samples were pre-incubated for 10 minutes at 4 °C. Reaction was initiated by the addition of enzyme to samples, which were then incubated for 30 minutes at 37 °C. The fluorescence of AMC (7-amino-4-methylcoumarin) released was monitored at 465 nm (excitation wavelength 360 nm), using fluorescence spectrophotometer.

15

The concentrations of the test compounds, which inhibit 50% of cathepsin s C, G, H, L and S activity, are represented as IC<sub>50</sub> in Table 2.

Table 2

Test		Cathepsins (IC <sub>50</sub> , nM)				
Compound	C	G	H	L	S	
5	Example 1	382	>240	>240	138	133
	Example 2	>1000	>1000	>1000	2100	2100
	Example 44	394	>690	>690	574	460
	Example 57	>1000	>1000	>1000	285	731
	Example 60	>1000	>1000	>1000	103	>150
10	Example 61	>1000	>1000	>1000	>300	>300
	Example 62	>1000	>1000	>1000	>450	>450
	Example 75	>1000	>1000	>1000	32.5	>150
	Example 76	>2460	>2460	>2460	971	204
	Example 80	>1000	>1000	>1000	381	109
15	Example 83	>810	>810	>810	352	177
	Example 84	>1000	>1000	>1000	986	212
	Example 94	>1000	>1000	>1000	1452	197
	Example 124	>1000	>1000	>1000	69.6	124
	Example 147	>1000	>1000	>1000	249	214
20	Example 148	>1000	>1000	>1000	178	141

As shown in Table 2, the compounds of the present invention have also a high selective inhibition effect against cathepsin K activity.


### Test 3: Pharmacokinetic Study

The compounds of the present invention were orally administered to rat with 10 mg/kg dose. The pharmacokinetic data were as follows:  $C_{\max}$  of 500 ~ 4,000 ng/ml, AUC of 1,500 ~ 6,500 ng-hr/ml, and the half-life ( $T_{1/2}$ ) of about 1 ~ 6 hours. The bioavailabilities of the compounds of the present invention by oral  
5 administrations, which were calculated after intravenous administrations of the same doses, were about 30% ~ 90%.

10

What is claimed is:


1. A compound of formula (I) or a pharmaceutically acceptable salt thereof:




wherein:

$R^1$  is a  $C_{1-6}$  alkyl group optionally substituted with phenyl,  $C_{1-6}$  alkoxy, or benzyloxy; a  $C_{2-6}$  alkenyl group optionally substituted with phenyl; a  $C_{3-6}$  cycloalkyl group; a  $C_{1-5}$  alkoxy group; a phenyl group substituted with halogen, phenyl, trifluoromethoxy, oxopyrrolidyl, mono- or di-  $C_{1-4}$  alkylamino or  $R^4$ - $C_{1-4}$  alkoxy (wherein,  $R^4$  is morpholine, pyrrolidine or piperidine); a furanyl group optionally substituted with one or more functional groups selected from the group consisting of  $C_{1-6}$  alkyl, halogen, and oxopyrrolidyl; a benzofuranyl group optionally substituted with  $C_{1-6}$  alkyl or  $R^4$ - $C_{1-4}$  alkoxy (wherein,  $R^4$  is morpholine, pyrrolidine or piperidine); a thiophenyl group substituted with  $C_{1-6}$  alkyl or halogen; a  $C_{1-6}$  alkyl-isoxazolyl group; a pyridyl group optionally substituted with halogen; a morpholinyl group; a benzothiophenyl group; a quinolinyl group; a pyrazinyl group; a benzyloxy group; an oxopyranyl group; a  $C_{1-6}$  alkyl-7H-imidazo[2,1-b]oxazolyl group; a  $C_{1-6}$  alkyl-chromon-2-yl group; or a (N-t-butoxycarbonyl)piperidinyl group, and


$R^2$  and  $R^3$  are, each independently, hydrogen; hydroxy; nitro; halogen; cyano; a  $C_{1-6}$  alkyl group optionally substituted with one or more halogen atoms;  $C_{1-5}$  alkoxy;  $C_{1-5}$  alkyl-thio; furyl; 1H-tetrazol-5-yl; oxazolyl; or a group selected from the formula consisting of (II), (III) and (IV):



(II)



(III)



(IV)

wherein,

R<sup>5</sup> is hydrogen; hydroxy; C<sub>1-6</sub> alkyl; C<sub>1-5</sub> alkoxy; mono- or di- C<sub>1-6</sub> alkylamino; or C<sub>3-6</sub> cycloalkylamino,

5 R<sup>6</sup> is C<sub>1-6</sub> alkyl; phenyl optionally substituted with a C<sub>1-4</sub> alkoxy group; benzyl optionally substituted with a C<sub>1-4</sub> alkoxy group,

R<sup>7</sup> and R<sup>8</sup> are, each independently, hydrogen; a C<sub>1-6</sub> alkylcarbonyl group optionally substituted with halogen, C<sub>1-4</sub> alkoxy, or phenyl; C<sub>2-6</sub> alkenylcarbonyl; C<sub>1-4</sub> alkoxy carbonyl; C<sub>3-6</sub> cycloalkylcarbonyl; benzoyl  
10 optionally substituted with one or more halogen atoms; mono- or di- C<sub>1-4</sub> alkylcarbonyl; or C<sub>1-4</sub> alkylsulfonyl, or bonded each ether to form a morpholine, azetidin-2-one, 3,3-dimethylazetidin-2-one, pyrrolidin-2-one, pyrrole, 2,5-dihydropyrrole, piperidin-2-one, oxazolidin-2-one, imidazolidin-2-one, imidazolidin-2,5-dione, tetrazole,  
15 1,1-dioxoisothiazolidine, or C<sub>1-6</sub> alkyl-aziridin-2-one ring, and

n is 0, 1, or 2.

2. The compound of formula (I) or its pharmaceutically acceptable salt of claim 1, wherein R<sup>1</sup> is a furanyl group optionally substituted with C<sub>1-6</sub> alkyl,  
20 halogen, or oxopyrrolidyl; or a benzofuranyl group optionally substituted with C<sub>1-6</sub> alkyl or R<sup>4</sup>-C<sub>1-4</sub> alkoxy (wherein, R<sup>4</sup> is morpholine, pyrrolidine or piperidine) and R<sup>2</sup> and R<sup>3</sup> are, each independently, hydrogen; halogen; or a group of formula (III) or (IV).

25 3. The compound of formula (I) or its pharmaceutically acceptable salt thereof of claim 1, which is selected from the group consisting of:

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
30 1-(2-nitrophenyl)piperidin-3-one;

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(4-fluoro-2-nitrophenyl)piperidin-3-one;

- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(5-chloro-2-nitrophenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(4-methanesulfonylphenyl)piperidin-3-one;
- 5 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-acetylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(4-cyanophenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
10 1-(2-cyano-3-fluorophenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-cyano-6-fluorophenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-cyano-5-fluorophenyl)piperidin-3-one;
- 15 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(3-cyano-5-fluorophenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-ethanesulfonylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
20 1-(4-acetylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(4-formylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-formylphenyl)piperidin-3-one;
- 25 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(4-nitrophenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(3-methyl-4-nitrophenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
30 1-(3-fluoro-2-formylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(3-chloro-2-formylphenyl)piperidin-3-one;

- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-formyl-4-nitrophenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-formyl-6-fluorophenyl)piperidin-3-one;
- 5 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-fluoro-3-formylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-isopropylsulfonylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
10 1-(4-fluoro-2-trifluoromethylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-cyanophenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-(4-morpholino)phenyl]piperidin-3-one;
- 15 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-ethoxyphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-fluoro-4-trifluoromethylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
20 1-(2-chloro-4-trifluoromethylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2,4-dichlorophenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(4-chloro-2-fluorophenyl)piperidin-3-one;
- 25 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-methoxy-5-cyanophenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(4-cyano-2-fluorophenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
30 1-[2-(2-furyl)phenyl]piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-bromophenyl)piperidin-3-one;



4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-methylthiophenyl)piperidin-3-one;

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-(tert-butoxycarbonyl)phenyl]piperidin-3-one;

5 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-ethoxycarbonylphenyl)piperidin-3-one;

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-methoxyphenyl)piperidin-3-one;

10 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(4-fluoro-2-methylphenyl)piperidin-3-one;

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-isopropoxyphenyl)piperidin-3-one;

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-ethylthiophenyl)piperidin-3-one;

15 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-(3,3-dimethylazetidin-2-one-1-yl)phenyl]piperidin-3-one;

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-(pyrrolidin-2-one-1-yl)phenyl]piperidin-3-one;

20 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-(piperidin-2-one-1-yl)phenyl]piperidin-3-one;

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-(2-oxazolidinon-1-yl)phenyl]piperidin-3-one;

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-(2-imidazolidinon-1-yl)phenyl]piperidin-3-one;

25 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-[N-(ethoxycarbonyl)amino]phenyl]piperidin-3-one;

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-[N-(methoxycarbonyl)amino]phenyl]piperidin-3-one;

30 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-[N-(4-chlorobutyryl)amino]phenyl]piperidin-3-one;

4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-[N-(isobutyryl)amino]phenyl]piperidin-3-one;

- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-[N-(methoxyacetyl)amino]phenyl]piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-[N-(cyclopropanecarbonyl)amino]phenyl]piperidin-3-one;
- 5 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-[N-(3-phenylpropionyl)amino]phenyl]piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-[N-(2-fluorobenzoyl)amino]phenyl]piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
10 1-[2-[N-(acetyl)amino]phenyl]piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-(2,5-dihydropyrrol-1-yl)phenyl]piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-(tetrazol-1-yl)phenyl]piperidin-3-one;
- 15 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-(pyrrol-1-yl)-5-fluorophenyl]piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-(pyrrol-1-yl)phenyl]piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
20 1-[2-(N,N-dimethylcarbamoyl)phenyl]piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-(N-methylcarbamoyl)phenyl]piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(2-carboxylphenyl)piperidin-3-one;
- 25 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-(4-carboxyphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-(1,3-oxazol-5-yl)phenyl]piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
30 1-(2-hydroxyphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-  
1-[2-(1H-tetrazol-5-yl)phenyl]piperidin-3-one;

- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methylsulfinylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-ethylsulfinylphenyl)piperidin-3-one;
- 5 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-isopropylsulfinylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-N-(acryloyl)aminophenyl]piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(3,3-dimethylureido)phenyl]piperidin-3-one;
- 10 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(methoxycarbonyl)phenyl]piperidin-3-one;
- 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-phenylpiperidin-3-one;
- 15 4-[N-[1-[N-(benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[2-(1,1-dioxo-isothiazolin-2-yl)phenyl]piperidin-3-one;
- 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(3-fluoro-2-methanesulfonylphenyl)piperidin-3-one;
- 20 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(4-fluoro-2-nitrophenyl)piperidin-3-one;
- 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(4-fluoro-2-formylphenyl)piperidin-3-one;
- 25 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-isopropylsulfonylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-phenylsulfonylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-(4-methoxybenzylsulfonylphenyl)piperidin-3-one;
- 30 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(4-fluoro-2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(4-methoxy-2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(4-fluoro-2-thiomethoxyphenyl)piperidin-3-one;

5 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-cyanophenyl)piperidin-3-one;

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(4-fluoro-2-methoxyacetaminophenyl)piperidin-3-one;

10 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(4-fluoro-2-methoxycarbonylaminophenyl)piperidin-3-one;

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(4-fluoro-2-cyclopropylcarbonylaminophenyl)piperidin-3-one;

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(4-fluoro-2-isobutyrylaminophenyl)piperidin-3-one;

15 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[4-fluoro-2-(4-chlorobutyryl)aminophenyl]piperidin-3-one;

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[4-fluoro-2-(3,3-dimethylazetidin-2-one-1-yl)phenyl]piperidin-3-one;

20 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(3-methyl-aziridin-2-one-1-yl)phenyl]piperidin-3-one;

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(azetidin-2-one-1-yl)phenyl]piperidin-3-one;

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[4-fluoro-2-(pyrrolidin-2-one-1-yl)phenyl]piperidin-3-one;

25 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(1,3-imidazolidin-2,5-dione-1-yl)phenyl]piperidin-3-one;

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[4-fluoro-2-(1,1-dioxo-isothiazolidin-2-yl)phenyl]piperidin-3-one;

30 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[4-fluoro-2-(piperidin-2-one-1-yl)phenyl]piperidin-3-one;

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[4-fluoro-2-(pyrrol-1-yl)]piperidin-3-one;

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[4-fluoro-2-(oxazol-4-yl)phenyl]piperidin-3-one;

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[4-fluoro-2-(N-methylcarbamoyl)phenyl]piperidin-3-one;

5 4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[2-(N,N-dimethylcarbamoyl)phenyl]piperidin-3-one;

4-[N-[1-[N-(furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[2-(N-cyclopropylcarbamoyl)phenyl]piperidin-3-one;

10 4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-aminophenyl)piperidin-3-one;

4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-nitrophenyl)piperidin-3-one;

4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

15 4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-ethanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[4-fluoro-2-(3,3-dimethyl-azetidin-2-one-1-yl)phenyl]piperidin-3-one;

20 4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[4-fluoro-2-(pyrrolidin-2-one-1-yl)phenyl]piperidin-3-one;

4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[4-fluoro-2-(oxazolidin-2-one-3-yl)phenyl]piperidin-3-one;

4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[4-fluoro-2-(imidazolidin-2-one-1-yl)phenyl]piperidin-3-one;

25 4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[2-(3,3-dimethyl-azetidin-2-one-1-yl)phenyl]piperidin-3-one;

4-[N-[1-[N-(5-methyl-isoxazol-3-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[2-(pyrrolidin-2-one-1-yl)phenyl]piperidin-3-one;

30 4-[N-[1-[N-(5-chloro-thiophen-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(5-chloro-thiophen-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-ethanesulfonylphenyl)piperidin-3-one;

- 4-[N-[1-[N-(5-chloro-thiophen-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-isopropanesulfonylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(5-chloro-thiophen-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(5-fluoro-2-methanesulfonylphenyl)piperidin-3-one;
- 5 4-[N-[1-[N-(5-methyl-thiophen-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(5-methyl-thiophen-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-isopropanesulfonylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(5-methyl-thiophen-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(5-fluoro-2-methanesulfonylphenyl)piperidin-3-one;
- 10 4-[N-[1-[N-(3-methyl-thiophen-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(2-methyl-furan-3-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;
- 15 4-[N-[1-[N-(5-methyl-furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(5-bromo-furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonyl-4-methoxyphenyl)piperidin-3-one;
- 4-[N-[1-[N-(4,5-dimethyl-furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;
- 20 4-[N-[1-[N-(4,5-dimethyl-furan-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[2-(pyrrolidin-2-one-1-yl)phenyl]piperidin-3-one;
- 4-[N-[1-[N-(morpholin-4-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;
- 25 4-[N-[1-[N-(benzothiophen-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzothiophen-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-tert-butoxycarbonylphenyl)piperidin-3-one;
- 4-[N-[1-[N-(benzothiophen-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-chloro-5-trifluoromethylphenyl)piperidin-3-one;
- 30 4-[N-[1-[N-(benzothiophen-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(4-chloro-2-fluoro-phenyl)piperidin-3-one;

4-[N-[1-[N-(benzothiophen-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methylphenyl)piperidin-3-one;

4-[N-[1-[N-(3-methyl-benzofuran-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

5 4-[N-[1-[N-(3-methyl-benzofuran-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(5-fluoro-2-trifluoromethylphenyl)piperidin-3-one;

4-[N-[1-[N-(3-methyl-benzofuran-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(5-cyano-2-methoxyphenyl)piperidin-3-one;

10 4-[N-[1-[N-(quinolin-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(quinolin-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-[2-(pyrrolidin-2-one-1-yl)phenyl]piperidin-3-one;

4-[N-[1-[N-(quinolin-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-nitrophenyl)piperidin-3-one;

15 4-[N-[1-[N-(pyrazin-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(pyrazin-2-ylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one dihydrochloride;

20 4-[N-[1-[N-(isonicotinylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(6-chloronicotinylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(6-chloronicotinylcarbonyl)amino]cyclohexylcarbonyl]amino]-1-(4-fluoro-2-nitrophenyl)piperidin-3-one;

25 4-[N-[1-[N-(pyridin-2-carbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(pyridin-2-carbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one hydrochloride;

30 4-[N-[1-[N-(4-fluorobenzoyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(4-fluorobenzoyl)amino]cyclohexylcarbonyl]amino]-1-(4-fluoro-2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(4-fluorobenzoyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonyl-4-methoxyphenyl)piperidin-3-one;

4-[N-[1-[N-(4-fluorobenzoyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-trifluoromethylphenyl)piperidin-3-one;

5 4-[N-[1-[N-(4-fluorobenzoyl)amino]cyclohexanecarbonyl]amino]-1-[(2-(3-phenylpropionylamino)phenyl)piperidin-3-one;

4-[N-[1-[N-(4-fluorobenzoyl)amino]cyclohexanecarbonyl]amino]-1-[2-(cyclopropylamino)phenyl]piperidin-3-one;

10 4-[N-[1-[N-(4-fluorobenzoyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(2,3,4-trifluorobenzoylamino)phenyl]piperidin-3-one;

4-[N-[1-[N-(4-biphenylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(4-trifluoromethylbenzoyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

15 4-[N-[1-[N-(4-(2-morpholinethoxy)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(4-(2-pyrrolidinethoxy)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

20 4-[N-[1-[N-(4-(2-piperidinethoxy)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(4-(2-morpholinethoxy)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one hydrochloride;

4-[N-[1-[N-(4-(2-pyrrolidinethoxy)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one hydrochloride;

25 4-[N-[1-[N-(4-(2-piperidinethoxy)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one hydrochloride;

4-[N-[1-[N-(5-(2-morpholinethoxy)benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

30 4-[N-[1-[N-(5-(2-pyrrolidinethoxy)benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(5-(2-piperidinethoxy)benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;



4-[N-[1-[N-(5-(2-morpholinethoxy)benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one hydrochloride;

5 4-[N-[1-[N-(5-(2-pyrrolidinethoxy)benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one hydrochloride;

4-[N-[1-[N-(5-(2-piperidinethoxy)benzofuran-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one hydrochloride;

10 4-[N-[1-[N-(4-(2-oxopyrrolidine)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(4-(2-oxopyrrolidine)furan-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(4-(2-morpholinethoxy)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-methanesulfonylaminophenyl)piperidin-3-one;

15 4-[N-[1-[N-(4-(2-morpholinethoxy)benzoyl)amino]cyclohexanecarbonyl]amino]-1-[4-fluoro-2-(pyrrolidin-2-one-1-yl)phenyl]piperidin-3-one;

4-[N-[1-[N-(4-(dimethylamino)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-methanesulfonylphenyl)piperidin-3-one;

20 4-[N-[1-[N-(4-(diethylamino)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(4-(dimethylamino)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-methanesulfonylphenyl)piperidin-3-one hydrochloride;

4-[N-[1-[N-(4-(diethylamino)benzoyl)amino]cyclohexanecarbonyl]amino]-1-(4-fluoro-2-methanesulfonylphenyl)piperidin-3-one hydrochloride;

25 4-[N-[1-[N-(3-phenylpropionyl)amino]cyclohexylcarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(2-methylcinnamoyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

30 4-[N-[1-[N-(benzyloxycarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-ethylamino)piperidin-3-one;

4-[N-[1-[N-(benzyloxycarbonyl)amino]cyclohexylcarbonyl]amino]-1-(2-trifluoromethylphenyl)piperidin-3-one;

4-[N-[1-[N-(pyran-2-one-5-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-((6-methyl)-7H-imidazo[2,1-b]oxazol-5-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

5 4-[N-[1-[N-((6-methyl)-chromon-2-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(isobutyryl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

10 4-[N-[1-[N-(isovaleryl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(3,3-dimethylacryloyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(2-methoxyacetyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

15 4-[N-[1-[N-(cyclopropanecarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-(cyclobutanecarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

20 4-[N-[1-[N-(cyclopentanecarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;



4-[N-[1-[N-(cyclohexanecarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

4-[N-[1-[N-((3-isobutyloxy)propionyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one;

25 4-[N-[1-[N-((3-benzyloxy)propionyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one; and

4-[N-[1-[N-((N-t-butoxycarbonyl)piperidin-4-ylcarbonyl)amino]cyclohexanecarbonyl]amino]-1-(2-methanesulfonylphenyl)piperidin-3-one.

30 4. A process for preparing a compound of formula (I) or a pharmaceutically acceptable salt thereof, which comprises oxidizing a compound of formula (V) with an oxidizing agent:




wherein,  $\text{R}^1$ ,  $\text{R}^2$ , and  $\text{R}^3$  are the same as defined in claim 1.

5

5. The process of claim 4, wherein the oxidizing agent is a pyridine- $\text{SO}_3$  complex, oxaryl chloride-dimethylsulfoxide, or pyridinium chlorochromate.

10



6. A compound of formula (V) or a pharmaceutically acceptable salt thereof:



wherein,  $\text{R}^1$ ,  $\text{R}^2$ , and  $\text{R}^3$  are the same as defined in claim 1..

15

7. A process for preparing a compound of formula (V) or a pharmaceutically acceptable salt thereof, which comprises reacting a compound of formula (VI) with a compound of formula (VII):



wherein, R<sup>1</sup>, R<sup>2</sup>, and R<sup>3</sup> are the same as defined in claim 1; and X is halogen, nitro, C<sub>1-7</sub> alkylsulfonyl, or trifluorosulfonate.

5

8. The process of claim 7, wherein the reaction is carried out in the presence of a base and a palladium catalyst.

9. The process of claim 8, wherein the palladium catalyst is palladium diacetate (Pd(OAc)<sub>2</sub>) or tris(dibenzylideneacetone)dipalladium (Pd<sub>2</sub>(dba)<sub>2</sub>).

10. A pharmaceutical compositions for inhibiting cysteine proteases comprising a therapeutically effective amount of the 1-phenylpiperidin-3-one derivative or a pharmaceutically acceptable salt thereof of claim 1 and a pharmaceutically acceptable carrier.

# INTERNATIONAL SEARCH REPORT

Application No.

PCT/KR03/01502

## A. CLASSIFICATION OF SUBJECT MATTER

**IPC7 C07D 405/12**

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: C07D 405/12, C07D 401/04, C07D 401/10

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched  
Korean Patents and application for inventions since 1975

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
MEDLINE, NPS, PAJ, CA on line, STN on line

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 0038687 A (SMITHKLINE BEECHAM CORP.) 06 JUL 2000 see claims 1-38, 59-61	1-10
A	WO 9808802 A (SMITHKLINE BEECHAM CORP.) 05 MAR 1998 see claims 1-10, 13, 14	1-3, 10
A	KR 1077201 A (YUHAN CO., LTD.) 17 AUG 2001 see claims 1-4	1-3, 10
A	WO 0048993 A (NOVARTIS-ERFINDUNGEN VERWALT.) 24 AUG 2000 see claims 1-10	1-3, 10

☐ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family


Date of the actual completion of the international search

27 OCTOBER 2003 (27.10.2003)

Date of mailing of the international search report

27 OCTOBER 2003 (27.10.2003)

Name and mailing address of the ISA/KR

 Korean Intellectual Property Office  
920 Dunsan-dong, Seo-gu, Daejeon 302-701,  
Republic of Korea

Facsimile No. 82-42-472-3556

Authorized officer

BAIK, Kyong UP

Telephone No. 82-42-481-5600



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/KR03/01502

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0038687 A	06.07.00	ZA 0104208 A	23.05.02
		US 20030144175 A	31.07.03
		US 20030044399 A	06.03.03
		US 20020147188 A	10.10.02
		TR 0101869 T	21.01.02
		PL 0350132 A	04.11.02
		NO 20013124 A	22.06.01
		JP 2002533397 T	08.10.02
		IL 0143142 A	21.04.02
		EP 1158986 A	05.12.01
		CZ 20012277 A	14.11.01
		CN 1350458 T	22.05.02
		CA 2356671 A	06.07.00
		BR 9916488 A	09.10.01
		AU 0019411 A	31.07.00
WO 9808802 A	05.03.98	JP 0517319 T	26.12.00
		EP 0923535 A	23.06.99
KR 1077201 A	17.08.01	None	
WO 0048993 A	24.08.00	GB 9903548 A	07.04.99
		AU 0028046 A	04.09.00